

Game Theory

3. Mixed Strategies

Albert-Ludwigs-Universität Freiburg



Bernhard Nebel and Robert Mattmüller
May 8, 2017

1 Mixed Strategies



- Definitions
- Support Lemma

Mixed Strategies
Definitions
Support Lemma
Nash's Theorem
Correlated Equilibria
Summary

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

3 / 56

Mixed Strategies



Mixed Strategies
Definitions
Support Lemma
Nash's Theorem
Correlated Equilibria
Summary

Observation: Not every strategic game has a pure-strategy Nash equilibrium (e. g. matching pennies).

Question:

- Can we do anything about that?
- Which strategy to play then?

Idea: Consider **randomized** strategies.

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

4 / 56

Mixed Strategies



Mixed Strategies
Definitions
Support Lemma
Nash's Theorem
Correlated Equilibria
Summary

Notation

Let X be a set.

Then $\Delta(X)$ denotes the set of **probability distributions** over X .

That is, each $p \in \Delta(X)$ is a mapping $p : X \rightarrow [0, 1]$ with

$$\sum_{x \in X} p(x) = 1.$$

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

5 / 56

A mixed strategy is a strategy where a player is allowed to randomize his action (throw a dice mentally and then act according to what he has decided to do for each outcome).

Definition (Mixed strategy)

Let $G = \langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ be a strategic game.

A **mixed strategy** of player i in G is a probability distribution $\alpha_i \in \Delta(A_i)$ over player i 's actions.

For $a_i \in A_i$, $\alpha_i(a_i)$ is the probability for playing a_i .

Terminology: When we talk about strategies in A_i specifically, to distinguish them from mixed strategies, we sometimes also call them **pure strategies**.

Definition (Mixed strategy profile)

A profile $\alpha = (\alpha_i)_{i \in N} \in \prod_{i \in N} \Delta(A_i)$ of mixed strategies induces a probability distribution p_α over $A = \prod_{i \in N} A_i$ as follows:

$$p_\alpha(a) = \prod_{i \in N} \alpha_i(a_i).$$

For $A' \subseteq A$, we define

$$p_\alpha(A') = \sum_{a \in A'} p_\alpha(a) = \sum_{a \in A'} \prod_{i \in N} \alpha_i(a_i).$$

Notation

Since each pure strategy $a_i \in A_i$ is equivalent to its induced mixed strategy \hat{a}_i

$$\hat{a}_i(a'_i) = \begin{cases} 1 & \text{if } a'_i = a_i \\ 0 & \text{otherwise,} \end{cases}$$

we sometimes abuse notation and write a_i instead of \hat{a}_i .

Example (Mixed strategies for matching pennies)

	H	T
H	1, -1	-1, 1
T	-1, 1	1, -1

$\alpha = (\alpha_1, \alpha_2)$, $\alpha_1(H) = 2/3$, $\alpha_1(T) = 1/3$, $\alpha_2(H) = 1/3$, $\alpha_2(T) = 2/3$.

This induces a probability distribution over $\{H, T\} \times \{H, T\}$:

$$\begin{aligned} p_\alpha(H, H) &= \alpha_1(H) \cdot \alpha_2(H) = 2/9, & u_1(H, H) &= +1, \\ p_\alpha(H, T) &= \alpha_1(H) \cdot \alpha_2(T) = 4/9, & u_1(H, T) &= -1, \\ p_\alpha(T, H) &= \alpha_1(T) \cdot \alpha_2(H) = 1/9, & u_1(T, H) &= -1, \\ p_\alpha(T, T) &= \alpha_1(T) \cdot \alpha_2(T) = 2/9, & u_1(T, T) &= +1. \end{aligned}$$

Definition (Expected utility)

Let $\alpha \in \prod_{i \in N} \Delta(A_i)$ be a mixed strategy profile.

The **expected utility** of α for player i is

$$U_i(\alpha) = U_i((\alpha_j)_{j \in N}) := \sum_{a \in A} p_\alpha(a) u_i(a) = \sum_{a \in A} \left(\prod_{j \in N} \alpha_j(a_j) \right) u_i(a).$$

Example (Mixed strategies for matching pennies (ctd.))

The expected utilities for player 1 and player 2 are

$$U_1(\alpha_1, \alpha_2) = -1/9 \quad \text{and} \quad U_2(\alpha_1, \alpha_2) = +1/9.$$

Remark: The expected utility functions U_i are linear in all mixed strategies.

Proposition

Let $\alpha \in \prod_{i \in N} \Delta(A_i)$ be a mixed strategy profile, $\beta_i, \gamma_i \in \Delta(A_i)$ mixed strategies, and $\lambda \in [0, 1]$. Then

$$U_i(\alpha_{-i}, \lambda \beta_i + (1 - \lambda) \gamma_i) = \lambda U_i(\alpha_{-i}, \beta_i) + (1 - \lambda) U_i(\alpha_{-i}, \gamma_i).$$

Moreover,

$$U_i(\alpha) = \sum_{a_i \in A_i} \alpha_i(a_i) \cdot U_i(\alpha_{-i}, a_i)$$

Proof.

Homework. □

Definition (Mixed extension)

Let $G = \langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ be a strategic game.

The **mixed extension** of G is the game $\langle N, (\Delta(A_i))_{i \in N}, (U_i)_{i \in N} \rangle$ where

- $\Delta(A_i)$ is the set of probability distributions over A_i and
- $U_i : \prod_{j \in N} \Delta(A_j) \rightarrow \mathbb{R}$ assigns to each mixed strategy profile α the expected utility for player i according to the induced probability distribution p_α .

Definition (Nash equilibrium in mixed strategies)

Let G be a strategic game.

A **Nash equilibrium in mixed strategies** (or **mixed-strategy Nash equilibrium**) of G is a Nash equilibrium in the mixed extension of G .

Intuition:

- It does not make sense to assign **positive probability** to a pure strategy that is **not a best response** to what the other players do.
- **Claim:** A profile of mixed strategies α is a Nash equilibrium if and only if everyone only plays best pure responses to what the others play.

Mixed
Strategies
Definitions
Support Lemma
Nash's
Theorem
Correlated
Equilibria
Summary

Definition (Support)

Let α_i be a mixed strategy.

The **support** of α_i is the set

$$\text{supp}(\alpha_i) = \{a_i \in A_i \mid \alpha_i(a_i) > 0\}$$

of actions played with nonzero probability.

Lemma (Support lemma)

Let $G = \langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ be a finite strategic game.

Then $\alpha^* \in \prod_{i \in N} \Delta(A_i)$ is a mixed-strategy Nash equilibrium in G if and only if for every player $i \in N$, every pure strategy in the support of α_i^* is a best response to α_{-i}^* .

For a single player—given all other players stick to their mixed strategies—it does not make a difference whether he plays the mixed strategy or whether he plays any single pure strategy from the support of the mixed strategy.

Mixed
Strategies
Definitions
Support Lemma
Nash's
Theorem
Correlated
Equilibria
Summary

Example (Support lemma)

Matching pennies, strategy profile $\alpha = (\alpha_1, \alpha_2)$ with

$$\alpha_1(H) = 2/3, \quad \alpha_1(T) = 1/3, \quad \alpha_2(H) = 1/3, \quad \text{and} \quad \alpha_2(T) = 2/3.$$

For α to be a Nash equilibrium, both actions in $\text{supp}(\alpha_2) = \{H, T\}$ have to be best responses to α_1 . Are they?

$$\begin{aligned} U_2(\alpha_1, H) &= \alpha_1(H) \cdot u_2(H, H) + \alpha_1(T) \cdot u_2(T, H) \\ &= 2/3 \cdot (-1) + 1/3 \cdot (+1) = -1/3, \end{aligned}$$

$$\begin{aligned} U_2(\alpha_1, T) &= \alpha_1(H) \cdot u_2(H, T) + \alpha_1(T) \cdot u_2(T, T) \\ &= 2/3 \cdot (+1) + 1/3 \cdot (-1) = 1/3. \end{aligned}$$

\Rightarrow Support lemma \Rightarrow $H \in \text{supp}(\alpha_2)$, but $H \notin B_2(\alpha_1)$.
 α can **not** be a Nash equilibrium.

Mixed
Strategies
Definitions
Support Lemma
Nash's
Theorem
Correlated
Equilibria
Summary

Proof.

\Rightarrow : Let α^* be a Nash equilibrium with $a_i \in \text{supp}(\alpha_i^*)$.

Assume that a_i is not a best response to α_{-i}^* . Because U_i is linear, player i can improve his utility by shifting probability in α_i^* from a_i to a better response.

This makes the modified α_i^* a better response than the original α_i^* , i. e., the original α_i^* was not a best response, which contradicts the assumption that α^* is a Nash equilibrium.

Mixed
Strategies
Definitions
Support Lemma
Nash's
Theorem
Correlated
Equilibria
Summary

Proof (ctd.)

“ \Leftarrow ”: Assume that α^* is not a Nash equilibrium.

Then there must be a player $i \in N$ and a strategy α'_i such that $U_i(\alpha^*_{-i}, \alpha'_i) > U_i(\alpha^*_{-i}, \alpha^*_i)$.

Because U_i is linear, there must be a pure strategy $a'_i \in \text{supp}(\alpha'_i)$ that has higher utility than some pure strategy $a''_i \in \text{supp}(\alpha^*_i)$.

Therefore, $\text{supp}(\alpha^*_i)$ does not only contain best responses to α^*_{-i} . \square

Example (Mixed-strategy Nash equilibria in BoS)

	B	S
B	2, 1	0, 0
S	0, 0	1, 2

We already know: (B, B) and (S, S) are pure Nash equilibria.

Possible supports (excluding “pure-vs-pure” strategies) are:

$$\{B\} \text{ vs. } \{B, S\}, \quad \{S\} \text{ vs. } \{B, S\}, \quad \{B, S\} \text{ vs. } \{B\}, \\ \{B, S\} \text{ vs. } \{S\} \quad \text{and} \quad \{B, S\} \text{ vs. } \{B, S\}$$

Observation: In Bach or Stravinsky, pure strategies have unique best responses. Therefore, there can be no Nash equilibria of “pure-vs-strictly-mixed” type.

Example (Mixed-strategy Nash equilibria in BoS (ctd.))

Consequence: Only need to search for additional Nash equilibria with support sets $\{B, S\}$ vs. $\{B, S\}$.

Assume that (α^*_1, α^*_2) is a Nash equilibrium with $0 < \alpha^*_1(B) < 1$ and $0 < \alpha^*_2(B) < 1$. Then

$$\begin{aligned} U_1(B, \alpha^*_2) &= U_1(S, \alpha^*_2) \\ \Rightarrow 2 \cdot \alpha^*_2(B) + 0 \cdot \alpha^*_2(S) &= 0 \cdot \alpha^*_2(B) + 1 \cdot \alpha^*_2(S) \\ \Rightarrow 2 \cdot \alpha^*_2(B) &= 1 - \alpha^*_2(B) \\ \Rightarrow 3 \cdot \alpha^*_2(B) &= 1 \\ \Rightarrow \alpha^*_2(B) &= 1/3 \quad (\text{and } \alpha^*_2(S) = 2/3) \end{aligned}$$

Similarly, we get $\alpha^*_1(B) = 2/3$ and $\alpha^*_1(S) = 1/3$.
The payoff profile of this equilibrium is $(2/3, 2/3)$.

Remark

Let $G = \langle \{1, 2\}, (A_i), (u_i) \rangle$ with $A_1 = \{T, B\}$ and $A_2 = \{L, R\}$ be a two-player game with two actions each, and (T, α^*_2) with $0 < \alpha^*_2(L) < 1$ be a Nash equilibrium of G .

Then at least one of the profiles (T, L) and (T, R) is also a Nash equilibrium of G .

Reason: Both L and R are best responses to T . Assume that T was neither a best response to L nor to R . Then B would be a better response than T both to L and to R .

With the linearity of U_1 , B would also be a better response to α^*_2 than T is. Contradiction.

Example

Consider the Nash equilibrium $\alpha^* = (\alpha_1^*, \alpha_2^*)$ with

$$\alpha_1^*(T) = 1, \quad \alpha_1^*(B) = 0, \quad \alpha_2^*(L) = 1/10, \quad \alpha_2^*(R) = 9/10$$

in the following game:

	L	R
T	1, 1	1, 1
B	2, 2	-5, -5

Here, (T, R) is also a Nash equilibrium.

- Definitions
- Kakutani's Fixpoint Theorem
- Proof of Nash's Theorem

Motivation: When does a strategic game have a mixed-strategy Nash equilibrium?

In the previous chapter, we discussed necessary and sufficient conditions for the existence of Nash equilibria for the special case of zero-sum games. Can we make other claims?

Theorem (Nash's theorem)

Every finite strategic game has a mixed-strategy Nash equilibrium.

Proof sketch.

Consider the set-valued function of best responses $B : \mathbb{R}^{\sum_i |A_i|} \rightarrow 2^{\mathbb{R}^{\sum_i |A_i|}}$ with

$$B(\alpha) = \prod_{i \in N} B_i(\alpha_{-i}).$$

A mixed strategy profile α is a fixed point of B if and only if $\alpha \in B(\alpha)$ if and only if α is a mixed-strategy Nash equilibrium.

The graph of B has to be connected. Then there is at least one point on the fixpoint diagonal. □

Nash's Theorem

Outline for the formal proof:

- 1 Review of necessary **mathematical definitions**
 \rightsquigarrow Subsection "Definitions"
- 2 **Statement of a fixpoint theorem** used to prove Nash's theorem (without proof)
 \rightsquigarrow Subsection "Kakutani's Fixpoint Theorem"
- 3 **Proof of Nash's theorem** using fixpoint theorem
 \rightsquigarrow Subsection "Proof of Nash's Theorem"

Nash's Theorem


Definitions

Definition


A set $X \subseteq \mathbb{R}^n$ is **closed** if X contains all its limit points, i. e., if $(x_k)_{k \in \mathbb{N}}$ is a sequence of elements in X and $\lim_{k \rightarrow \infty} x_k = x$, then also $x \in X$.

Example

Closed:



Not closed:



Nash's Theorem

Definitions


Definition

A set $X \subseteq \mathbb{R}^n$ is **bounded** if for each $i = 1, \dots, n$ there are lower and upper bounds $a_i, b_i \in \mathbb{R}$ such that


$$X \subseteq \prod_{i=1}^n [a_i, b_i].$$

Example

Bounded:



Not bounded:



Nash's Theorem

Definitions


Definition

A set $X \subseteq \mathbb{R}^n$ is **convex** if for all $x, y \in X$ and all $\lambda \in [0, 1]$,


$$\lambda x + (1 - \lambda)y \in X.$$

Example

Convex:



Not convex:



Nash's Theorem

Definitions



Mixed
Strategies

Nash's
Theorem

Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem

Correlated
Equilibria

Summary

Definition

For a function $f : X \rightarrow 2^X$, the **graph** of f is the set

$$\text{Graph}(f) = \{(x, y) \mid x \in X, y \in f(x)\}.$$

Nash's Theorem

Kakutani's Fixpoint Theorem



Mixed
Strategies

Nash's
Theorem

Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem

Correlated
Equilibria

Summary

Theorem (Kakutani's fixpoint theorem)

Let $X \subseteq \mathbb{R}^n$ be a nonempty, closed, bounded and convex set and let $f : X \rightarrow 2^X$ be a function such that

- for all $x \in X$, the set $f(x) \subseteq X$ is nonempty and convex, and
- $\text{Graph}(f)$ is closed.

Then there is an $x \in X$ with $x \in f(x)$, i. e., f has a fixpoint.

Proof.

See Shizuo Kakutani, A generalization of Brouwer's fixed point theorem, 1941, or your favorite advanced calculus textbook, or the Internet.

For German speakers: Harro Heuser, Lehrbuch der Analysis, Teil 2, also has a proof (Abschnitt 232). □

Nash's Theorem

Kakutani's Fixpoint Theorem



Mixed
Strategies

Nash's
Theorem

Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem


Correlated
Equilibria

Summary


Example

Let $X = [0, 1]$.

Kakutani's theorem applicable:



Kakutani's theorem not applicable:



Nash's Theorem

Proof



Mixed
Strategies

Nash's
Theorem

Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem

Correlated
Equilibria

Summary

Proof.

Apply Kakutani's fixpoint theorem using $X = \mathcal{A} = \prod_{i \in N} \Delta(A_i)$ and $f = B$, where $B(\alpha) = \prod_{i \in N} B_i(\alpha_{-i})$.

We have to show:

- 1 \mathcal{A} is nonempty,
- 2 \mathcal{A} is closed,
- 3 \mathcal{A} is bounded,
- 4 \mathcal{A} is convex,
- 5 $B(\alpha)$ is nonempty for all $\alpha \in \mathcal{A}$,
- 6 $B(\alpha)$ is convex for all $\alpha \in \mathcal{A}$, and
- 7 $\text{Graph}(B)$ is closed.

Nash's Theorem

Proof



Proof (ctd.)

Some notation:

- Assume without loss of generality that $N = \{1, \dots, n\}$.
- A profile of mixed strategies can be written as a vector of $M = \sum_{i \in N} |A_i|$ real numbers in the interval $[0, 1]$ such that numbers for the same player add up to 1.

For example, $\alpha = (\alpha_1, \alpha_2)$ with $\alpha_1(T) = 0.7$, $\alpha_1(M) = 0.0$, $\alpha_1(B) = 0.3$, $\alpha_2(L) = 0.4$, $\alpha_2(R) = 0.6$ can be seen as the vector

$$\underbrace{(0.7, 0.0, 0.3)}_{\alpha_1}, \underbrace{(0.4, 0.6)}_{\alpha_2}$$

- This allows us to interpret the set \mathcal{A} of mixed strategy profiles as a subset of \mathbb{R}^M .

Mixed Strategies

Nash's Theorem
Definitions
Kakutani's Fixpoint Theorem
Proof of Nash's Theorem

Correlated Equilibria

Summary

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

35 / 56

Nash's Theorem

Proof



Proof (ctd.)

- \mathcal{A} nonempty: Trivial. \mathcal{A} contains the tuple

$$(1, \underbrace{0, \dots, 0}_{|A_1| - 1 \text{ times}}, \dots, 1, \underbrace{0, \dots, 0}_{|A_n| - 1 \text{ times}}).$$

- \mathcal{A} closed: Let $\alpha_1, \alpha_2, \dots$ be a sequence in \mathcal{A} that converges to $\lim_{k \rightarrow \infty} \alpha_k = \alpha$. Suppose $\alpha \notin \mathcal{A}$. Then either there is some component of α that is less than zero or greater than one, or the components for some player i add up to a value other than one.

Since α is a limit point, the same must hold for some α_k in the sequence. But then, $\alpha_k \notin \mathcal{A}$, a contradiction. Hence \mathcal{A} is closed.

Mixed Strategies

Nash's Theorem
Definitions
Kakutani's Fixpoint Theorem
Proof of Nash's Theorem

Correlated Equilibria

Summary

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

36 / 56

Nash's Theorem

Proof



Proof (ctd.)

- \mathcal{A} bounded: Trivial. All entries are between 0 and 1, i.e., \mathcal{A} is bounded by $[0, 1]^M$.
- \mathcal{A} convex: Let $\alpha, \beta \in \mathcal{A}$ and $\lambda \in [0, 1]$, and consider $\gamma = \lambda \alpha + (1 - \lambda) \beta$. Then

$$\begin{aligned} \min(\gamma) &= \min(\lambda \alpha + (1 - \lambda) \beta) \\ &\geq \lambda \cdot \min(\alpha) + (1 - \lambda) \cdot \min(\beta) \\ &\geq \lambda \cdot 0 + (1 - \lambda) \cdot 0 = 0, \end{aligned}$$

and similarly, $\max(\gamma) \leq 1$.

Hence, all entries in γ are still in $[0, 1]$.

Mixed Strategies

Nash's Theorem
Definitions
Kakutani's Fixpoint Theorem
Proof of Nash's Theorem

Correlated Equilibria

Summary

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

37 / 56

Nash's Theorem

Proof



Proof (ctd.)

- \mathcal{A} convex (ctd.): Let $\tilde{\alpha}, \tilde{\beta}$ and $\tilde{\gamma}$ be the sections of α, β and γ , respectively, that determine the probability distribution for player i . Then

$$\begin{aligned} \sum \tilde{\gamma} &= \sum (\lambda \tilde{\alpha} + (1 - \lambda) \tilde{\beta}) \\ &= \lambda \cdot \sum \tilde{\alpha} + (1 - \lambda) \cdot \sum \tilde{\beta} \\ &= \lambda \cdot 1 + (1 - \lambda) \cdot 1 = 1. \end{aligned}$$

Hence, all probabilities for player i in γ still sum up to 1.

Altogether, $\gamma \in \mathcal{A}$, and therefore, \mathcal{A} is convex.

Mixed Strategies

Nash's Theorem
Definitions
Kakutani's Fixpoint Theorem
Proof of Nash's Theorem

Correlated Equilibria

Summary

May 8, 2017

B. Nebel, R. Mattmüller – Game Theory

38 / 56

Nash's Theorem

Proof



Proof (ctd.)

- 5 **$B(\alpha)$ nonempty:** For a fixed α_{-i} , U_i is linear in the mixed strategies of player i , i. e., for $\beta_i, \gamma_i \in \Delta(A_i)$,

$$U_i(\alpha_{-i}, \lambda\beta_i + (1-\lambda)\gamma_i) = \lambda U_i(\alpha_{-i}, \beta_i) + (1-\lambda)U_i(\alpha_{-i}, \gamma_i) \quad (1)$$

for all $\lambda \in [0, 1]$.

Hence, U_i is continuous on $\Delta(A_i)$.

Continuous functions on closed and bounded sets take their maximum in that set.

Therefore, $B_i(\alpha_{-i}) \neq \emptyset$ for all $i \in N$, and thus $B(\alpha) \neq \emptyset$.

Mixed
Strategies
Nash's
Theorem
Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem
Correlated
Equilibria
Summary

Nash's Theorem

Proof



Proof (ctd.)

- 6 **$B(\alpha)$ convex:** This follows, since each $B_i(\alpha_{-i})$ is convex. To see this, let $\alpha'_i, \alpha''_i \in B_i(\alpha_{-i})$.

Then $U_i(\alpha_{-i}, \alpha'_i) = U_i(\alpha_{-i}, \alpha''_i)$.

With Equation (1), this implies

$$\lambda\alpha'_i + (1-\lambda)\alpha''_i \in B_i(\alpha_{-i}).$$

Hence, $B_i(\alpha_{-i})$ is convex.

- 7 **$Graph(B)$ closed:** Let (α^k, β^k) be a convergent sequence in $Graph(B)$ with $\lim_{k \rightarrow \infty} (\alpha^k, \beta^k) = (\alpha, \beta)$.

So, $\alpha^k, \beta^k, \alpha, \beta \in \prod_{i \in N} \Delta(A_i)$ and $\beta^k \in B(\alpha^k)$.

We need to show that $(\alpha, \beta) \in Graph(B)$, i. e., that $\beta \in B(\alpha)$.

Mixed
Strategies
Nash's
Theorem
Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem
Correlated
Equilibria
Summary

Nash's Theorem

Proof



Proof (ctd.)

- 7 **$Graph(B)$ closed (ctd.):** It holds for all $i \in N$:

$$\begin{aligned} U_i(\alpha_{-i}, \beta_i) &\stackrel{(D)}{=} U_i(\lim_{k \rightarrow \infty} (\alpha_{-i}^k, \beta_i^k)) \\ &\stackrel{(C)}{=} \lim_{k \rightarrow \infty} U_i(\alpha_{-i}^k, \beta_i^k) \\ &\stackrel{(B)}{\geq} \lim_{k \rightarrow \infty} U_i(\alpha_{-i}^k, \beta'_i) \quad \text{for all } \beta'_i \in \Delta(A_i) \\ &\stackrel{(C)}{=} U_i(\lim_{k \rightarrow \infty} \alpha_{-i}^k, \beta'_i) \quad \text{for all } \beta'_i \in \Delta(A_i) \\ &\stackrel{(D)}{=} U_i(\alpha_{-i}, \beta'_i) \quad \text{for all } \beta'_i \in \Delta(A_i). \end{aligned}$$

(D): def. α_i, β_i ; (C) continuity; (B) β_i^k best response to α_{-i}^k .

Mixed
Strategies
Nash's
Theorem
Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem
Correlated
Equilibria
Summary

Nash's Theorem

Proof



Proof (ctd.)

- 7 **$Graph(B)$ closed (ctd.):** It follows that β_i is a best response to α_{-i} for all $i \in N$.

Thus, $\beta \in B(\alpha)$ and finally $(\alpha, \beta) \in Graph(B)$.

Therefore, all requirements of Kakutani's fixpoint theorem are satisfied.

Applying Kakutani's theorem establishes the existence of a fixpoint of B , which is, by definition/construction, the same as a mixed-strategy Nash equilibrium. \square

Mixed
Strategies
Nash's
Theorem
Definitions
Kakutani's Fixpoint
Theorem
Proof of Nash's
Theorem
Correlated
Equilibria
Summary

3 Correlated Equilibria

Correlated Equilibria

Recall: There are three Nash equilibria in Bach or Stravinsky

- (B, B) with payoff profile $(2, 1)$
- (S, S) with payoff profile $(1, 2)$
- (α_1^*, α_2^*) with payoff profile $(2/3, 2/3)$ where
 - $\alpha_1^*(B) = 2/3, \alpha_1^*(S) = 1/3,$
 - $\alpha_2^*(B) = 1/3, \alpha_2^*(S) = 2/3.$

Idea: Use a publicly visible coin toss to decide which action from a mixed strategy is played. This can lead to higher payoffs.

Correlated Equilibria

Example (Correlated equilibrium in BoS)

With a **fair coin** that both players can observe, the players can agree to play as follows:

- If the coin shows heads, both play B .
- If the coin shows tails, both play S .

This is **stable** in the sense that no player has an incentive to deviate from this agreed-upon rule, as long as the other player keeps playing his/her strategy (cf. definition of Nash equilibria).

Expected payoffs: $(3/2, 3/2)$ instead of $(2/3, 2/3)$.

Observations and Information Partitions

We assume that observations are made based on a finite probability space (Ω, π) , where Ω is a set of **states** and π is a **probability measure** on Ω .

Agents might not be able to distinguish all states from each other. In order to model this, we assume for each player i an **information partition** $\mathcal{P}_i = \{P_{i1}, P_{i2}, \dots, P_{ik}\}$. This means that $\bigcup_{j=1}^{j=ik} P_j = \Omega$ and for all $P_i, P_k \in \mathcal{P}_i$ with $P_j \neq P_k$, we have $P_j \cap P_k = \emptyset$.

Example: $\Omega = \{x, y, z\}$, $\mathcal{P}_1 = \{\{x\}, \{y, z\}\}$, $\mathcal{P}_2 = \{\{x, y\}, \{z\}\}$.

We say that a function $f: \Omega \rightarrow X$ **respects an information partition** for player i if $f(\omega) = f(\omega')$ whenever $\omega \in P_i$ and $\omega' \in P_i$ for some $P_i \in \mathcal{P}_i$.

Example: f respects \mathcal{P}_1 if $f(y) = f(z)$.

Definition

A **correlated equilibrium of a strategic game** $\langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ consists of

- a finite probability space (Ω, π) ,
- for each player $i \in N$ an **information partition** \mathcal{P}_i of Ω ,
- for each player $i \in N$ a function $\sigma_i : \Omega \rightarrow A_i$ that respects \mathcal{P}_i (σ_i is player i 's **strategy**)

such that for every $i \in N$ and every function $\tau_i : \Omega \rightarrow A_i$ that respects \mathcal{P}_i (i.e. for every possible strategy of player i) we have

$$\sum_{\omega \in \Omega} \pi(\omega) u_i(\sigma_{-i}(\omega), \sigma_i(\omega)) \geq \sum_{\omega \in \Omega} \pi(\omega) u_i(\sigma_{-i}(\omega), \tau_i(\omega)). \quad (2)$$

	L	R
T	6, 6	2, 7
B	7, 2	0, 0

Equilibria: (T, R) with $(2, 7)$, (B, L) with $(7, 2)$, and mixed $((\frac{2}{3}, \frac{1}{3}), (\frac{2}{3}, \frac{1}{3}))$ with $(4\frac{2}{3}, 4\frac{2}{3})$.

Assume $\Omega = \{x, y, z\}$, $\pi(x) = \frac{1}{3}$, $\pi(y) = \frac{1}{3}$, $\pi(z) = \frac{1}{3}$.

Assume further $\mathcal{P}_1 = \{\{x\}, \{y, z\}\}$, $\mathcal{P}_2 = \{\{x, y\}, \{z\}\}$.

Set $\sigma_1(x) = B$, $\sigma_1(y) = \sigma_1(z) = T$ and $\sigma_2(x) = \sigma_2(y) = L$, $\sigma_2(z) = R$.

Then both player play optimally and get a payoff profile of $(5, 5)$.

Proposition

For every mixed strategy Nash equilibrium α of a finite strategic game $\langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$, there is a correlated equilibrium $\langle (\Omega, \pi), (\mathcal{P}_i), (\sigma_i) \rangle$ in which for each player i the distribution on A_i induced by σ_i is α_i .

This means that correlated equilibria are a generalization of Nash equilibria.

Proof.

Let $\Omega = A$ and define $\pi(a) = \prod_{i \in N} \alpha_i(a_i)$. For each player i , let $a \in P$ and $b \in P$ for $P \in \mathcal{P}_i$ if $a_i = b_i$. Define $\sigma_i(a) = a_i$ for each $a \in A$.

Then $\langle (\Omega, \pi), (\mathcal{P}_i), (\sigma_i) \rangle$ is a correlated equilibrium since the left hand side of (2) is the Nash equilibrium payoff and for each player i at least as good any other strategy τ_i respecting the information partition. Further, the distribution induced by σ_i is α_i . □

Proposition

Let $G = \langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ be a strategic game. Any convex combination of correlated equilibrium payoff profiles of G is a correlated equilibrium payoff profile of G .

Proof idea: From given equilibria and weighting factors, create a new one by combining them orthogonally, using the weighting factors.

Proof

Proof.

Let u^1, \dots, u^K be the payoff profiles and let $(\lambda^1, \dots, \lambda^K) \in \mathbb{R}^K$ with $\lambda^l \geq 0$ and $\sum_{l=1}^K \lambda^l = 1$. For each l let $(\Omega^l, \pi^l), (\mathcal{P}_i^l), (\sigma_i^l)$

be a correlated equilibrium generating payoff u^l . Wlog. assume all Ω^l 's are disjoint.

Now we define a correlated equilibrium generating the payoff $\sum_{l=1}^K \lambda^l u^l$. Let $\Omega = \bigcup_l \Omega^l$. For any $\omega \in \Omega$ define $\pi(\omega) = \lambda^l \pi^l(\omega)$ where l is such that $\omega \in \Omega^l$. For each $i \in N$ let $\mathcal{P}_i = \bigcup_l \mathcal{P}_i^l$ and set $\sigma_i(\omega) = \sigma_i^l(\omega)$ where l is such that $\omega \in \Omega^l$.

□

Basically, first throw a dice for which CE to go for, then proceed in this CE.

4 Summary

Summary

- **Mixed strategies** allow randomization.
- **Characterization** of mixed-strategy Nash equilibria: players only play best responses with positive probability (**support lemma**).
- **Nash's Theorem**: Every finite strategic game has a mixed-strategy Nash equilibrium.
- **Correlated equilibria** can lead to higher payoffs.
- All Nash equilibria are correlated equilibria, but not *vice versa*.