

Game Theory

4. Algorithms

Albert-Ludwigs-Universität Freiburg



Bernhard Nebel and Robert Mattmüller

May 15th, 2017

1 Motivation



Motivation

Linear Programming

Zero-Sum Games

General Finite Two-Player Games

Summary

May 15th, 2017

B. Nebel, R. Mattmüller – Game Theory

3 / 36

Motivation



- **We know:** In finite strategic games, mixed-strategy Nash equilibria are guaranteed to **exist**.
- **We don't know:** How to systematically **find them**?
- **Challenge:** There are **infinitely many** mixed strategy profiles to consider. How to do this in finite time?

This chapter:

- Computation of mixed-strategy Nash equilibria for **finite zero-sum games**.
- Computation of mixed-strategy Nash equilibria for **general finite two player games**.

Motivation

Linear Programming

Zero-Sum Games

General Finite Two-Player Games

Summary

May 15th, 2017

B. Nebel, R. Mattmüller – Game Theory

4 / 36

2 Linear Programming



Motivation

Linear Programming

Zero-Sum Games

General Finite Two-Player Games

Summary

May 15th, 2017

B. Nebel, R. Mattmüller – Game Theory

6 / 36

Digression:

We briefly discuss **linear programming** because we will use this technique to find Nash equilibria.

Goal of linear programming:

Solving a system of linear inequalities over n real-valued variables while **optimizing** some **linear objective function**.

Example

Production of two sorts of items with time requirements and profit per item. Objective: Maximize profit.

	Cutting	Assembly	Postproc.	Profit per item
(x) sort 1	25	60	68	30
(y) sort 2	75	60	34	40
per day	≤ 450	≤ 480	≤ 476	maximize!


Goal: Find numbers of pieces x of sort 1 and y of sort 2 to be produced per day such that the resource constraints are met and the objective function is maximized.

Example (ctd., formalization)

$$\begin{aligned}
 &x \geq 0, y \geq 0 & (1) \\
 &25x + 75y \leq 450 \quad (\text{or } y \leq 6 - \frac{1}{3}x) & (2) \\
 &60x + 60y \leq 480 \quad (\text{or } y \leq 8 - x) & (3) \\
 &68x + 34y \leq 476 \quad (\text{or } y \leq 14 - 2x) & (4) \\
 &\text{maximize } z = 30x + 40y & (5)
 \end{aligned}$$

- Inequalities (1)–(4): Admissible solutions
(They form a **convex set** in $\mathbb{R}^2$.)
- Line (5): Objective function

Example (ctd., visualization)



Definition (Linear program)

A **linear program** (LP) in standard form consists of

- n real-valued variables x_i ; n coefficients b_i ;
- m constants c_j ; $n \cdot m$ coefficients a_{ij} ;
- m constraints of the form

$$c_j \leq \sum_{i=1}^n a_{ij} x_i,$$

- and an objective function to be minimized ($x_i \geq 0$)

$$\sum_{i=1}^n b_i x_i.$$

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Solution of an LP:

assignment of values to the x_i **satisfying the constraints** and **minimizing the objective function**.

Remarks:

- **Maximization instead of minimization:** easy, just change the signs of all the b_i 's, $i = 1, \dots, n$.
- **Equalities** instead of inequalities: $x + y \leq c$ if and only if there is a $z \geq 0$ such that $x + y + z = c$ (z is called a **slack variable**).

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Solution algorithms:

- Usually, one uses the **simplex algorithm** (which is worst-case exponential!).
- There are also polynomial-time algorithms such as interior-point or ellipsoid algorithms.

Tools and libraries:

- **lp_solve**
- CLP
- GLPK
- CPLEX
- gurobi

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Mixed-Strategy Nash Equilibria in Finite Zero-Sum Games



Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

We start with **finite zero-sum games** for two reasons:

- They are **easier to solve** than general finite two-player games.
- Understanding how to solve finite zero-sum games **facilitates understanding** how to solve general finite two-player games.

Mixed-Strategy Nash Equilibria in Finite Zero-Sum Games



Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

In the following, we will **exploit the zero-sum property** of a game G when searching for mixed-strategy Nash equilibria. For that, we need the following result.

Proposition

Let G be a finite zero-sum game. Then the mixed extension of G is also a zero-sum game.

Proof.

Homework. □

Mixed-Strategy Nash Equilibria in Finite Zero-Sum Games



Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Let G be a finite zero-sum game with mixed extension G' .

Then we know the following:

- 1 **Previous proposition implies:** G' is also a zero-sum game.
- 2 **Nash's theorem implies:** G' has a Nash equilibrium.
- 3 **Maximinimizer theorem + (1) + (2) implies:** Nash equilibria and pairs of maximinimizers in G' are the same.

Mixed-Strategy Nash Equilibria in Finite Zero-Sum Games



Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Consequence:

When **looking for mixed-strategy Nash equilibria in G** , it is sufficient to **look for pairs of maximinimizers in G'** .

Method: Linear Programming

Approach:

- Let $G = \langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ be a finite zero-sum game:
 - $N = \{1, 2\}$.
 - A_1 and A_2 are finite.
 - $U_1(\alpha, \beta) = -U_2(\alpha, \beta)$ for all $\alpha \in \Delta(A_1), \beta \in \Delta(A_2)$.
- Player 1 looks for a maximinimizer mixed strategy α .
- For each possible α of player 1:
 - Determine expected utility against best response of pl. 2. (Only need to consider **finitely many pure** candidates for best responses because of Support Lemma).
 - Maximize expected utility over all possible α .

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

- **Result:** maximinimizer α for player 1 in G' (= Nash equilibrium strategy for player 1)
- **Analogously:** obtain maximinimizer β for player 2 in G' (= Nash equilibrium strategy for player 2)
- **With maximinimizer theorem:** we can combine α and β into a **Nash equilibrium**.

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

“For each possible α of player 1, determine expected utility against best response of player 2, and maximize.”

translates to the following LP:

$$\begin{aligned} \alpha(a) &\geq 0 \quad \text{for all } a \in A_1 \\ \sum_{a \in A_1} \alpha(a) &= 1 \\ U_1(\alpha, b) = \sum_{a \in A_1} \alpha(a) \cdot u_1(a, b) &\geq u \quad \text{for all } b \in A_2 \\ \text{Maximize } u. \end{aligned}$$

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

Example (Matching pennies)

	H	T
H	1, -1	-1, 1
T	-1, 1	1, -1

Linear program for player 1:

Maximize u subject to the constraints

$$\begin{aligned} \alpha(H) &\geq 0, \alpha(T) \geq 0, \alpha(H) + \alpha(T) = 1, \\ \alpha(H) \cdot u_1(H, H) + \alpha(T) \cdot u_1(T, H) &= \alpha(H) - \alpha(T) \geq u, \\ \alpha(H) \cdot u_1(H, T) + \alpha(T) \cdot u_1(T, T) &= -\alpha(H) + \alpha(T) \geq u. \end{aligned}$$

Solution: $\alpha(H) = \alpha(T) = 1/2, u = 0$.

Motivation
Linear Programming
Zero-Sum Games
General Finite Two-Player Games
Summary

- **Remark:** There is an alternative encoding based on the observation that in zero-sum games that have a Nash equilibrium, maximinimization and minimaximization yield the same result.
- **Idea:** Formulate linear program with inequalities

$$U_1(a, \beta) \leq u \quad \text{for all } a \in A_1$$

and minimize u . Analogously for β .

- For general finite two-player games, the **LP approach does not work**.
- Instead, use instances of the **linear complementarity problem (LCP)**:
 - Linear (in-)equalities as with LPs.
 - Additional constraints of the form $x_i \cdot y_i = 0$ (or equivalently $x_i = 0 \vee y_i = 0$) for variables $X = \{x_1, \dots, x_k\}$ and $Y = \{y_1, \dots, y_k\}$, and $i \in \{1, \dots, k\}$.
 - **no** objective function.
- With LCPs, we can compute Nash equilibria for **arbitrary** finite two-player games.

Let A_1 and A_2 be finite and let (α, β) be a Nash equilibrium with payoff profile (u, v) . Then consider this LCP encoding:

$$u - U_1(a, \beta) \geq 0 \quad \text{for all } a \in A_1 \quad (6)$$

$$v - U_2(\alpha, b) \geq 0 \quad \text{for all } b \in A_2 \quad (7)$$

$$\alpha(a) \cdot (u - U_1(a, \beta)) = 0 \quad \text{for all } a \in A_1 \quad (8)$$

$$\beta(b) \cdot (v - U_2(\alpha, b)) = 0 \quad \text{for all } b \in A_2 \quad (9)$$

$$\alpha(a) \geq 0 \quad \text{for all } a \in A_1 \quad (10)$$

$$\sum_{a \in A_1} \alpha(a) = 1 \quad (11)$$

$$\beta(b) \geq 0 \quad \text{for all } b \in A_2 \quad (12)$$

$$\sum_{b \in A_2} \beta(b) = 1 \quad (13)$$

Remarks about the encoding:

- In (8) and (9): for instance,

$$\alpha(a) \cdot (u - U_1(a, \beta)) = 0$$

if and only if

$$\alpha(a) = 0 \quad \text{or} \quad u - U_1(a, \beta) = 0.$$

This holds in every Nash equilibrium, because:

- if $a \notin \text{supp}(\alpha)$, then $\alpha(a) = 0$, and
- if $a \in \text{supp}(\alpha)$, then $a \in B_1(\beta)$, thus $U_1(a, \beta) = u$.
- With additional variables, the above LCP formulation can be transformed into LCP normal form.

Motivation
Linear Pro-
gramming
Zero-Sum
Games
General
Finite
Two-Player
Games
Summary

Theorem

A mixed strategy profile (α, β) with payoff profile (u, v) is a Nash equilibrium if and only if it is a solution to the LCP encoding over (α, β) and (u, v) .

Proof.

- **Nash equilibria are solutions to the LCP:** Obvious because of the support lemma.
- **Solutions to the LCP are Nash equilibria:** Let (α, β, u, v) be a solution to the LCP. Because of (10)–(13), α and β are mixed strategies.

Motivation
Linear Pro-
gramming
Zero-Sum
Games
General
Finite
Two-Player
Games
Summary

Proof (ctd.)

- **Solutions to the LCP are Nash equilibria (ctd.):** Because of (6), u is at least the maximal payoff over all possible pure responses, and because of (8), u is exactly the maximal payoff.

If $\alpha(a) > 0$, then, because of (8), the payoff for player 1 against β is u .

The linearity of the expected utility implies that α is a best response to β .

Analogously, we can show that β is a best response to α and hence (α, β) is a Nash equilibrium with payoff profile (u, v) .

□

Motivation
Linear Pro-
gramming
Zero-Sum
Games
General
Finite
Two-Player
Games
Summary

Naïve algorithm:

Enumerate all $(2^n - 1) \cdot (2^m - 1)$ possible pairs of support sets.

For each such pair $(\text{supp}(\alpha), \text{supp}(\beta))$:

- Convert the LCP into an LP:
 - Linear (in-)equalities are preserved.
 - Constraints of the form $\alpha(a) \cdot (u - U_1(a, \beta)) = 0$ are replaced by a new linear equality:
 - $u - U_1(a, \beta) = 0$, if $a \in \text{supp}(\alpha)$, and
 - $\alpha(a) = 0$, otherwise,
 - Analogously for $\beta(b) \cdot (v - U_2(\alpha, b)) = 0$.
 - Objective function: maximize constant zero function.
- Apply solution algorithm for LPs to the transformed program.

Motivation
Linear Pro-
gramming
Zero-Sum
Games
General
Finite
Two-Player
Games
Summary

- Runtime of the naïve algorithm: $O(p(n+m) \cdot 2^{n+m})$, where p is some polynomial.
- Better in practice: Lemke-Howson algorithm.
- Complexity:
 - unknown whether $\text{LCP SOLVE} \in \mathbf{P}$.
 - $\text{LCP SOLVE} \in \mathbf{NP}$ is clear (naïve algorithm can be seen as a nondeterministic polynomial-time algorithm).

- Computation of mixed-strategy Nash equilibria for **finite zero-sum games** using **linear programs**.
→ polynomial-time computation
- Computation of mixed-strategy Nash equilibria for **general finite two player games** using **linear complementarity problem**.
→ computation in **NP**.