Multi-Agent Systems

B. Nebel, F. Lindner, T. Engesser Summer Semester 2017 University of Freiburg Department of Computer Science


Exercise Sheet 10 Due: July 24th, 2017, 10:00

Exercise 10.1 (The Shapley Value, 3+2)

- (a) Prove that the Shapley value satisfies the axioms Symmetry, Dummy player, and Additivity.
- (b) Prove that the Shapley value is *feasible* and *efficient*.

Exercise 10.2 (Taxi Cost Sharing, 2+2+2+2+2)

Consider a scenario for sharing taxis and taxi fares that is slightly more complex than the one presented in the lecture. There are four agents who each want to be transported from a common starting point (a taxi stand) to their individual destinations (A, B, C, and D). The costs a taxi charges for a route can be calculated by adding up the costs for the single route segments, which are given as edge labels in the following graph:



E.g., the route Start-A-C has a cost of 9 + 7 = 16. Since taxis take always the shortest and cheapest route, the taxi fare for a coalition is always the cost of the cheapest route starting from "Start" and containing all the destinations of the agents in the coalition.

- (a) Model the problem as cooperative game (N, v).
- (b) Calculate the Shapley value $\Psi(N, v)$.
- (c) Find the socially optimal coalition structure CS^* for (N, v).
- (d) Calculate the Shapley value $\Psi(N', v)$ for each coalition $N' \in CS^*$.
- (e) Which of the Shapley values $\Psi(X, v)$ calculated in (b) and (d) are in the core of their respective games (X, v)? Prove your answer!