

Game Theory

6. Extensive Games

Albert-Ludwigs-Universität Freiburg



**UNI
FREIBURG**

Bernhard Nebel and Robert Mattmüller

May 9th, 2018

1 Motivation



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

- **So far:** All players move **simultaneously**, and then the outcome is determined.
- **Often in practice:** Several moves in **sequence** (e. g. in chess).
↪ cannot be directly reflected by strategic games.
- **Extensive games** (with perfect information) reflect such situations by modeling games as **game trees**.
- **Idea:** Players have several decision points where they can decide how to play.
- **Strategies:** Mappings from decision points in the game tree to actions to be played.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

2 Definitions



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition (Extensive game with perfect information)

An **extensive game with perfect information** is a tuple

$\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ that consists of:

- A finite non-empty set N of **players**.
- A set H of (finite or infinite) sequences, called **histories**, such that
 - the empty sequence $\langle \rangle \in H$,
 - H is **closed under prefixes**: if $\langle a^1, \dots, a^k \rangle \in H$ for some $k \in \mathbb{N} \cup \{\infty\}$, and $l < k$, then also $\langle a^1, \dots, a^l \rangle \in H$, and
 - H is **closed under limits**: if for some infinite sequence $\langle a^i \rangle_{i=1}^\infty$, we have $\langle a^i \rangle_{i=1}^k \in H$ for all $k \in \mathbb{N}$, then $\langle a^i \rangle_{i=1}^\infty \in H$.

All infinite histories and all histories $\langle a^i \rangle_{i=1}^k \in H$, for which there is no a^{k+1} such that $\langle a^i \rangle_{i=1}^{k+1} \in H$ are called **terminal histories** Z . Components of a history are called **actions**.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary



Definition (Extensive game with perfect information, ctd.)

- A **player function** $P : H \setminus Z \rightarrow N$ that determines which player's turn it is to move after a given nonterminal history.
- For each player $i \in N$, a **utility function** (or **payoff function**) $u_i : Z \rightarrow \mathbb{R}$ defined on the set of terminal histories.

The game is called **finite**, if H is finite. It has a **finite horizon**, if the length of histories is bounded from above.

Assumption: All ingredients of Γ are **common knowledge** amongst the players of the game.

Terminology: In the following, we will simply write **extensive games** instead of **extensive games with perfect information**.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property


Kuhn's
Theorem

Two
Extensions

Summary

Example (Division game)

- Two identical objects should be **divided** among two players.
- **Player 1 proposes** an allocation.
- **Player 2 agrees or rejects**.
 - **On agreement:** Allocation as proposed.
 - **On rejection:** Nobody gets anything.



Motivation

Definitions

Solution
Concepts


One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Example (Division game, formally)



- $N = \{1, 2\}$
- $H = \{\langle \rangle, \langle (2,0) \rangle, \langle (1,1) \rangle, \langle (0,2) \rangle, \langle (2,0), y \rangle, \langle (2,0), n \rangle, \dots\}$
- $P(\langle \rangle) = 1, P(h) = 2$ for all $h \in H \setminus Z$ with $h \neq \langle \rangle$
- $u_1(\langle (2,0), y \rangle) = 2, u_2(\langle (2,0), y \rangle) = 0$, etc.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Notation:

Let $h = \langle a^1, \dots, a^k \rangle$ be a history, and a an action.

- Then (h, a) is the history $\langle a^1, \dots, a^k, a \rangle$.
- If $h' = \langle b^1, \dots, b^\ell \rangle$, then (h, h') is the history $\langle a^1, \dots, a^k, b^1, \dots, b^\ell \rangle$.
- The set of actions from which player $P(h)$ can choose after a history $h \in H \setminus Z$ is written as

$$A(h) = \{a \mid (h, a) \in H\}.$$

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition (Strategy in an extensive game)

A **strategy** of a player i in an extensive game $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ is a function s_i that assigns to each nonterminal history $h \in H \setminus Z$ with $P(h) = i$ an action $a \in A(h)$. The set of strategies of player i is denoted as S_i .

Remark: Strategies require us to assign actions to histories h , even if it is clear that they will never be played (e. g., because h will never be reached because of some earlier action).

Notation (for finite games): A strategy for a player is written as a string of actions at decision nodes as visited in a breadth-first order.

Motivation

Definitions

Solution
Concepts


One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Example (Strategies in an extensive game)



- Strategies for player 1: AE , AF , BE and BF
- Strategies for player 2: C and D .

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions


Summary

Definition (Outcome)

The **outcome** $O(s)$ of a strategy profile $s = (s_i)_{i \in N}$ is the (possibly infinite) terminal history $h = \langle a^i \rangle_{i=1}^k$, with $k \in \mathbb{N} \cup \{\infty\}$, such that for all $\ell \in \mathbb{N}$ with $0 \leq \ell < k$,

$$s_{P(\langle a^1, \dots, a^\ell \rangle)}(\langle a^1, \dots, a^\ell \rangle) = a^{\ell+1}.$$

Example (Outcome)



$$O(AF, C) = \langle A, C, F \rangle$$

$$O(AE, D) = \langle A, D \rangle.$$

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

3 Solution Concepts



Motivation

Definitions

**Solution
Concepts**

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition (Nash equilibrium in an extensive game)

A **Nash equilibrium** in an extensive game $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ is a strategy profile s^* such that for every player $i \in N$ and for all strategies $s_i \in S_i$,

$$u_i(O(s_{-i}^*, s_i^*)) \geq u_i(O(s_{-i}^*, s_i)).$$

Definition (Induced strategic game)

The strategic game G **induced** by an extensive game

$\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ is defined by $G = \langle N, (A'_i)_{i \in N}, (u'_i)_{i \in N} \rangle$,
where

- $A'_i = S_i$ for all $i \in N$, and
- $u'_i(a) = u_i(O(a))$ for all $i \in N$.

Proposition

The Nash equilibria of an extensive game Γ are exactly the Nash equilibria of the induced strategic game G of Γ . □

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Remarks:

- Each extensive game can be transformed into a strategic game, but the resulting game can be exponentially larger.
- The other direction does not work, because in extensive games, we do not have simultaneous actions.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property


Kuhn's
Theorem

Two
Extensions

Summary

Example (Empty threat)

Extensive game:



Strategic form:

	L	R
T	0, 0	2, 1
B	1, 2	1, 2

Nash equilibria: (B, L) and (T, R) .

However, (B, L) is not realistic:

- Player 1 plays B , “fearing” response L to T .
- But player 2 would never play L in the extensive game.
 $\rightsquigarrow (B, L)$ involves “empty threat”.

Strategies:

- Player 1: T and B
- Player 2: L and R

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Idea: Exclude empty threats.

How? Demand that a strategy profile is not only a Nash equilibrium in the strategic form, but also in every subgame.

Definition (Subgame)

A **subgame** of an extensive game $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$, starting after history h , is the game $\Gamma(h) = \langle N, H|_h, P|_h, (u_i|_h)_{i \in N} \rangle$, where

- $H|_h = \{h' \mid (h, h') \in H\}$,
- $P|_h(h') = P(h, h')$ for all $h' \in H|_h$, and
- $u_i|_h(h') = u_i(h, h')$ for all $h' \in H|_h$.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition (Strategy in a subgame)

Let Γ be an extensive game and $\Gamma(h)$ a subgame of Γ starting after some history h .

For each strategy s_i of Γ , let $s_i|_h$ be the strategy induced by s_i for $\Gamma(h)$. Formally, for all $h' \in H|_h$,

$$s_i|_h(h') = s_i(h, h').$$

The outcome function of $\Gamma(h)$ is denoted by O_h .



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition (Subgame-perfect equilibrium)


A strategy profile s^* in an extensive game $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ is a **subgame-perfect equilibrium** if and only if for every player $i \in N$ and every nonterminal history $h \in H \setminus Z$ with $P(h) = i$,

$$u_i|_h(O_h(s^*_{-i}|_h, s_i^*|_h)) \geq u_i|_h(O_h(s^*_{-i}|_h, s_i))$$


for every strategy $s_i \in S_i$ in subgame $\Gamma(h)$.

Two Nash equilibria:

- (T, R) : **subgame-perfect**, because:
 - In history $h = \langle T \rangle$: subgame-perfect.
 - In history $h = \langle \rangle$: player 1 obtains utility 1 when choosing B and utility of 2 when choosing T .
- (B, L) : **not subgame-perfect**, since L does not maximize the utility of player 2 in history $h = \langle T \rangle$.



Example (Subgame-perfect equilibria in division game)



Equilibria in subgames:

- in $\Gamma(\langle(2,0)\rangle)$: y and n
- in $\Gamma(\langle(1,1)\rangle)$: only y
- in $\Gamma(\langle(0,2)\rangle)$: only y
- in $\Gamma(\langle\rangle)$: $((2,0), yyy)$ and $((1,1), nyy)$

Nash equilibria (red: empty threat):

- $((2,0), yyy)$, $((2,0), yyn)$, $((2,0), yny)$, $((2,0), ynn)$,
 $((2,0), nny)$, $((2,0), nnn)$,
- $((1,1), nyy)$, $((1,1), nyn)$,
- $((0,2), nny)$, $((0,2), nnn)$.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

4 One-Deviation Property



Motivation

Definitions

Solution
Concepts

**One-
Deviation
Property**

Kuhn's
Theorem

Two
Extensions

Summary

- **Existence:**
 - Does every extensive game have a subgame-perfect equilibrium?
 - If not, which extensive games do have a subgame-perfect equilibrium?
- **Computation:**
 - If a subgame-perfect equilibrium exists, how to compute it?
 - How complex is that computation?

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Positive case (a subgame-perfect equilibrium exists):

- **Step 1:** Show that it suffices to consider **local** deviations from strategies (for finite-horizon games).
- **Step 2:** Show how to **systematically explore such local deviations** to find a subgame-perfect equilibrium (for finite games).

Step 1: One-Deviation Property



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Definition

Let Γ be a finite-horizon extensive game. Then $\ell(\Gamma)$ denotes the length of the longest history of Γ .

Step 1: One-Deviation Property



Definition (One-deviation property)

A strategy profile s^* in an extensive game $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ satisfies the one-deviation property if and only if for every player $i \in N$ and every nonterminal history $h \in H \setminus Z$ with $P(h) = i$,

$$u_i|_h(O_h(s_{-i}^*|_h, s_i^*|_h)) \geq u_i|_h(O_h(s_{-i}^*|_h, s_i))$$

for every strategy $s_i \in S_i$ in subgame $\Gamma(h)$ that differs from $s_i^*|_h$ only in the action it prescribes after the initial history of $\Gamma(h)$.

Note: Without the highlighted parts, this is just the definition of subgame-perfect equilibria!

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 1: One-Deviation Property



Lemma

Let $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ be a **finite-horizon** extensive game. Then a strategy profile s^* is a subgame-perfect equilibrium of Γ if and only if it satisfies the one-deviation property.

Proof

- (\Rightarrow) Clear.
- (\Leftarrow) By contradiction:

Suppose that s^* is not a subgame-perfect equilibrium.

Then there is a history h and a player i such that s_i is a profitable deviation for player i in subgame $\Gamma(h)$.

...

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary


Step 1: One-Deviation Property



Proof (ctd.)

- (\Leftarrow) ... WLOG, the number of histories h' with $s_i(h') \neq s_i^*|_h(h')$ is at most $\ell(\Gamma(h))$ and hence finite (finite horizon assumption!), since deviations not on resulting outcome path are irrelevant.

Illustration: strategies $s_1^*|_h = AGILN$ and $s_2^*|_h = CF$ red:



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions


Summary

Step 1: One-Deviation Property



Proof (ctd.)

- (\Leftarrow) ... Illustration for WLOG assumption: Assume $s_1 = BHKMO$ (blue) profitable deviation:



Then only B and O really matter.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions


Summary

Step 1: One-Deviation Property



Proof (ctd.)

- (\Leftarrow) ... Illustration for WLOG assumption: And hence $\tilde{s}_1 = BGILO$ (blue) also profitable deviation:



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 1: One-Deviation Property



Proof (ctd.)

■ (\Leftarrow) ...

Choose profitable deviation s_i in $\Gamma(h)$ with minimal number of deviation points (such s_i must exist).

Let h^* be the longest history in $\Gamma(h)$ with $s_i(h^*) \neq s_i^*|_h(h^*)$, i.e., “deepest” deviation point for s_i .

Then in $\Gamma(h, h^*)$, $s_i|_{h^*}$ differs from $s_i^*|_{(h, h^*)}$ only in the initial history.

Moreover, $s_i|_{h^*}$ is a profitable deviation in $\Gamma(h, h^*)$, since h^* is the *longest* history in $\Gamma(h)$ with $s_i(h^*) \neq s_i^*|_h(h^*)$.

So, $\Gamma(h, h^*)$ is the desired subgame where a one-step deviation is sufficient to improve utility. □

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property


Kuhn's
Theorem

Two
Extensions

Summary

Step 1: One-Deviation Property

Example



To show that (AHI, CE) is a subgame-perfect equilibrium, it suffices to check these deviating strategies:

Player 1:

- G in subgame $\Gamma(\langle A, C \rangle)$
- K in subgame $\Gamma(\langle B, F \rangle)$
- BHI in Γ

Player 2:

- D in subgame $\Gamma(\langle A \rangle)$
- F in subgame $\Gamma(\langle B \rangle)$

In particular, e.g., no need to check if strategy BGK of player 1 is profitable in Γ .

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary


Step 1: One-Deviation Property

Remark on Infinite-Horizon Games



The corresponding proposition for infinite-horizon games **does not hold**.

Counterexample (one-player case):



Strategy s_i with $s_i(h) = S$ for all $h \in H \setminus Z$

- satisfies one deviation property, but
- is not a subgame-perfect equilibrium, since it is dominated by s_i^* with $s_i^*(h) = C$ for all $h \in H \setminus Z$.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

5 Kuhn's Theorem



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

**Kuhn's
Theorem**

Two
Extensions

Summary

Step 2: Kuhn's Theorem



Theorem (Kuhn)

Every **finite** extensive game has a subgame-perfect equilibrium.

Proof idea:

- Proof is **constructive** and builds a subgame-perfect equilibrium bottom-up (aka **backward induction**).
- For those familiar with the Foundations of AI lecture: generalization of Minimax algorithm to general-sum games with possibly more than two players.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem


Two
Extensions

Summary

Step 2: Kuhn's Theorem



Example



$$s_2(\langle A \rangle) = C$$

$$t_1(\langle A \rangle) = 1$$

$$t_2(\langle A \rangle) = 5$$

$$s_2(\langle B \rangle) = F$$

$$t_1(\langle B \rangle) = 0$$

$$t_2(\langle B \rangle) = 8$$

$$s_1(\langle \rangle) = A$$

$$t_1(\langle \rangle) = 1$$

$$t_2(\langle \rangle) = 5$$

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 2: Kuhn's Theorem



A bit more formally:

Proof

Let $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$ be a finite extensive game.

Construct a subgame-perfect equilibrium by induction on $\ell(\Gamma(h))$ for all subgames $\Gamma(h)$. In parallel, construct functions $t_i : H \rightarrow \mathbb{R}$ for all players $i \in N$ s. t. $t_i(h)$ is the payoff for player i in a subgame-perfect equilibrium in subgame $\Gamma(h)$.

Base case: If $\ell(\Gamma(h)) = 0$, then $t_i(h) = u_i(h)$ for all $i \in N$.

...

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 2: Kuhn's Theorem



Proof (ctd.)

Inductive case: If $t_j(h)$ already defined for all $h \in H$ with $\ell(\Gamma(h)) \leq k$, consider $h^* \in H$ with $\ell(\Gamma(h^*)) = k + 1$ and $P(h^*) = i$.

For all $a \in A(h^*)$, $\ell(\Gamma(h^*, a)) \leq k$, let

$$s_i(h^*) := \operatorname{argmax}_{a \in A(h^*)} t_i(h^*, a) \quad \text{and}$$

$$t_j(h^*) := t_j(h^*, s_i(h^*)) \quad \text{for all players } j \in N.$$

Inductively, we obtain a strategy profile s that satisfies the one-deviation property.

With the one-deviation property lemma it follows that s is a subgame-perfect equilibrium. □

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 2: Kuhn's Theorem



- **In principle:** sample subgame-perfect equilibrium effectively computable using the technique from the above proof.
- **In practice:** often game trees not enumerated in advance, hence unavailable for backward induction.
- E.g., for branching factor b and depth m , procedure needs time $O(b^m)$.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 2: Kuhn's Theorem

Remark on Infinite Games



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Corresponding proposition for infinite games **does not hold**.

Counterexamples (both for one-player case):

A) finite horizon, infinite branching factor:

Infinitely many actions $a \in A = [0, 1)$ with payoffs $u_1(\langle a \rangle) = a$ for all $a \in A$.


There exists no subgame-perfect equilibrium in this game.

Step 2: Kuhn's Theorem

Remark on Infinite Games



B) infinite horizon, finite branching factor:



$$u_1(CCC\dots) = 0 \text{ and } u_1(\underbrace{CC\dots C}_n S) = n + 1.$$

No subgame-perfect equilibrium.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

Step 2: Kuhn's Theorem



Uniqueness:

Kuhn's theorem tells us nothing about uniqueness of subgame-perfect equilibria. However, if no two histories get the same evaluation by any player, then the subgame-perfect equilibrium is unique.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

**Kuhn's
Theorem**

Two
Extensions

Summary

Extended Example: Pirate Game



- 1 There are 5 *rational* pirates, A, B, C, D and E . They find 100 gold coins. They must decide how to distribute them.
- 2 The pirates have a strict order of *seniority*: A is senior to B , who is senior to C , who is senior to D , who is senior to E .
- 3 The pirate world's rules of distribution say that the most senior pirate first *proposes* a distribution of coins. The pirates, including the proposer, then *vote* on whether to accept this distribution (in order from most junior to senior). In case of a tie vote, the proposer has the casting vote. If the distribution is accepted, the coins are disbursed and the *game ends*. If not, the proposer is thrown overboard from the pirate ship and dies, and the next most senior pirate makes a new proposal to apply the method again.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary



- 4 The pirates do not trust each other, and will neither make nor honor any promises between pirates apart from a proposed distribution plan that gives a whole number of gold coins to each pirate.
- 5 Pirates base their decisions on three factors. First of all, each pirate wants to *survive*. Second, everything being equal, each pirate wants to *maximize the number of gold coins* each receives. Third, each pirate would prefer to *throw another overboard*, if all other results would otherwise be equal.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

- Players $N = \{A, B, C, D, E\}$;
- actions are:
 - proposals by a pirate: $\langle A : x_A, B : x_B, C : x_C, D : x_D, E : x_E \rangle$, with $\sum_{i \in \{A, B, C, D, E\}} x_i = 100$;
 - votings: y for accepting, n for rejecting;
- histories are sequences of a proposal, followed by votings of the alive pirates;
- utilities:
 - for pirates who are alive: utilities are according to the accepted proposal plus $x/100$, x being the number of dead pirates;
 - for dead pirates: -100.

Remark: Very large game tree!

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary



- 1 Assume only D and E are still alive. D can propose $\langle A : 0, B : 0, C : 0, D : 100, E : 0 \rangle$, because D has the casting vote!
- 2 Assume C , D , and E are alive. For C it is enough to offer 1 coin to E to get his vote: $\langle A : 0, B : 0, C : 99, D : 0, E : 1 \rangle$.
- 3 Assume B , C , D , and E are alive. B offering D one coin is enough because of the casting vote:
 $\langle A : 0, B : 99, C : 0, D : 1, E : 0 \rangle$.
- 4 Assume A , B , C , D , and E are alive. A offering C and E each one coin is enough: $\langle A : 98, B : 0, C : 1, D : 0, E : 1 \rangle$
(note that giving 1 to D instead to E does not help).

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary

6 Two Extensions

- Simultaneous Moves
- Chance

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

**Two
Extensions**

Simultaneous
Moves

Chance

Summary

Definition

An **extensive game with simultaneous moves** is a tuple $\Gamma = \langle N, H, P, (u_i)_{i \in N} \rangle$, where

- N, H, P and (u_i) are defined as before, and
- $P : H \rightarrow 2^N$ assigns to each nonterminal history a **set** of players to move; for all $h \in H \setminus Z$, there exists a family $(A_i(h))_{i \in P(h)}$ such that

$$A(h) = \{a \mid (h, a) \in H\} = \prod_{i \in P(h)} A_i(h).$$

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

- **Intended meaning of simultaneous moves:** All players from $P(h)$ move simultaneously.
- **Strategies:** Functions $s_i : h \mapsto a_i$ with $a_i \in A_i(h)$.
- **Histories:** Sequences of vectors of actions.
- **Outcome:** Terminal history reached when tracing strategy profile.
- **Payoffs:** Utilities at outcome history.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary



Remark:

- The **one-deviation property still holds** for extensive game with perfect information and simultaneous moves.
- **Kuhn's theorem does not hold** for extensive game with simultaneous moves.

Example: MATCHING PENNIES can be viewed as extensive game with simultaneous moves. No Nash equilibrium/subgame-perfect equilibrium.

		player 2	
		<i>H</i>	<i>T</i>
player 1	<i>H</i>	1, -1	-1, 1
	<i>T</i>	-1, 1	1, -1

⇒ Need more sophisticated solution concepts (cf. mixed strategies). Not covered in this lecture.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

Setting:

- Three players have to split a cake fairly.
- Player 1 suggest split: shares $x_1, x_2, x_3 \in [0, 1]$ s.t.
 $x_1 + x_2 + x_3 = 1$.
- Then players 2 and 3 **simultaneously** and **independently** decide whether to accept (“y”) or reject (“n”) the suggested splitting.
- If both accept, each player i gets his allotted share (utility x_i). Otherwise, no player gets anything (utility 0).

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

Simultaneous Moves

Example: Three-Person Cake Splitting Game



Formally:

$$N = \{1, 2, 3\}$$

$$X = \{(x_1, x_2, x_3) \in [0, 1]^3 \mid x_1 + x_2 + x_3 = 1\}$$

$$H = \{\langle \rangle\} \cup \{\langle x \rangle \mid x \in X\} \cup \{\langle x, z \rangle \mid x \in X, z \in \{y, n\} \times \{y, n\}\}$$

$$P(\langle \rangle) = \{1\}$$

$$P(\langle x \rangle) = \{2, 3\} \text{ for all } x \in X$$

$$u_i(\langle x, z \rangle) = \begin{cases} 0 & \text{if } z \in \{(y, n), (n, y), (n, n)\} \\ x_i & \text{if } z = (y, y). \end{cases} \text{ for all } i \in N$$

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

Subgame-perfect equilibria:

- **Subgames after legal split** (x_1, x_2, x_3) by player 1:
 - NE (y, y) (both accept)
 - NE (n, n) (neither accepts)
 - If $x_2 = 0$, NE (n, y) (only player 3 accepts)
 - If $x_3 = 0$, NE (y, n) (only player 2 accepts)

Subgame-perfect equilibria (ctd.):

■ Entire game:

Let s_2 and s_3 be any two strategies of players 2 and 3 such that for all splits $x \in X$ the profile $(s_2(\langle x \rangle), s_3(\langle x \rangle))$ is one of the NEs from above.

Let $X_y = \{x \in X \mid s_2(\langle x \rangle) = s_3(\langle x \rangle) = y\}$ be the set of splits accepted under s_2 and s_3 . Distinguish three cases:

- $X_y = \emptyset$ or $x_1 = 0$ for all $x \in X_y$. Then (s_1, s_2, s_3) is a subgame-perfect equilibrium for any possible s_1 .
- $X_y \neq \emptyset$ and there are splits $x_{\max} = (x_1, x_2, x_3) \in X_y$ that maximize $x_1 > 0$. Then (s_1, s_2, s_3) is a subgame-perfect equilibrium if and only if $s_1(\langle \cdot \rangle)$ is such a split x_{\max} .
- $X_y \neq \emptyset$ and there are no splits $(x_1, x_2, x_3) \in X_y$ that maximize x_1 . Then there is no subgame-perfect equilibrium, in which player 2 follows strategy s_2 and player 3 follows strategy s_3 .

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

Definition

An **extensive game with chance moves** is a tuple

$\Gamma = \langle N, H, P, f_c, (u_i)_{i \in N} \rangle$, where

- N, A, H and u_i are defined as before,
- the player function $P : H \setminus Z \rightarrow N \cup \{c\}$ can also take the value **c** for a **chance node**, and
- for each $h \in H \setminus Z$ with $P(h) = c$, the function $f_c(\cdot|h)$ is a probability distribution on $A(h)$ such that the probability distributions for all $h \in H$ are independent of each other.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves

Chance

Summary

- **Intended meaning of chance moves:** In chance node, an applicable action is chosen randomly with probability according to f_c .
- **Strategies:** Defined as before.
- **Outcome:** For a given strategy profile, the outcome is a probability distribution on the set of terminal histories.
- **Payoffs:** For player i , U_i is the expected payoff (with weights according to outcome probabilities).

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property


Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves
Chance

Summary

Example



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves

Chance

Summary



Remark:

The one-deviation property and Kuhn's theorem still hold in the presence of chance moves. When proving Kuhn's theorem, **expected** utilities have to be used.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Simultaneous
Moves

Chance

Summary

7 Summary



Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary



- For **finite-horizon extensive games**, it suffices to consider **local deviations** when looking for better strategies.
- For infinite-horizon games, this is not true in general.
- Every **finite extensive game** has a **subgame-perfect equilibrium**.
- This does not generally hold for infinite games, no matter if game is infinite due to infinite branching factor or infinitely long histories (or both).

- With **chance moves**, one deviation property and Kuhn's theorem still hold.
- With **simultaneous moves**, Kuhn's theorem no longer holds.

Motivation

Definitions

Solution
Concepts

One-
Deviation
Property

Kuhn's
Theorem

Two
Extensions

Summary