

Principles of Knowledge Representation and Reasoning

Nonmonotonic Reasoning III: Cumulative Logics

Albert-Ludwigs-Universität Freiburg



Bernhard Nebel, Stefan Wöfl, and Julien Hué
December 7 & 12 2012

1 Introduction



- Motivation
- Properties
- Derived Rules in **C**
- Undesirable Properties

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

December 7 & 12 2012

Nebel, Wöfl, Hué – KRR

3 / 59

Motivation



- Conventional NM logics are based on (ad hoc) modifications of the logical machinery (proofs/models).
- **Nonmonotonicity** is only a **negative** characterization:
From $\Theta \vdash \varphi$, it does not necessarily follow $\Theta \cup \{\psi\} \vdash \varphi$.
- Could we have a constructive **positive** characterization of default reasoning?

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

December 7 & 12 2012

Nebel, Wöfl, Hué – KRR

4 / 59

Plausible consequences



- In classical logics, we have the logical consequence relation $\alpha \models \beta$: If α is true, then also β is true.
- Instead, we will study the relation of **plausible consequence** $\alpha \vdash \beta$: If α is all we know, can we conclude β ?
- $\alpha \vdash \beta$ does not imply $\alpha \wedge \alpha' \vdash \beta$!
Compare to conditional probability: $P(\beta|\alpha) \neq P(\beta|\alpha, \alpha')$!
- Find rules that characterize $\vdash \dots$
For example: if $\alpha \vdash \beta$ and $\alpha \vdash \gamma$, then $\alpha \vdash \beta \wedge \gamma$.
- Write down all such rules \dots
- \dots and find a **semantic characterization** of \vdash !

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

December 7 & 12 2012

Nebel, Wöfl, Hué – KRR

5 / 59

Reflexivity (Ref):

$$\frac{}{\alpha \sim \alpha}$$

- **Rationale:** If α holds, this normally implies α .
- **Example:** Tom goes to a party normally implies that Tom goes to a party.

Let $\Delta = \langle D, W \rangle$ be a propositional default theory.
Define the relation \sim_{Δ} as follows:

$$\alpha \sim_{\Delta} \beta \iff \langle D, W \cup \{\alpha\} \rangle \vdash \beta$$

$\alpha \sim_{\Delta} \beta$ means that β is a skeptical conclusion of $\langle D, W \cup \{\alpha\} \rangle$.

Proposition

Default logic satisfies Reflexivity.

Proof.

The question is: does α follow skeptically from $\Delta' = \langle D, W \cup \{\alpha\} \rangle$?
For each extension E of Δ' , it holds $W \cup \{\alpha\} \subseteq E$ (by definition).
Hence $\alpha \in E$, and thus α belongs to all extensions of Δ' . \square

Left Logical Equivalence (LLE):

$$\frac{\models \alpha \leftrightarrow \beta, \alpha \sim \gamma}{\beta \sim \gamma}$$

- **Rationale:** It is not the syntactic form, but the content that is responsible for what we conclude normally.
- **Example:** Assume that Tom goes or Peter goes normally implies Mary goes. Then we would expect that Peter goes or Tom goes normally implies Mary goes.

Proposition

Default logic satisfies Left Logical Equivalence.

Proof.

Assume $\models \alpha \leftrightarrow \beta$ and $\alpha \sim_{\Delta} \gamma$ (with $\Delta = \langle D, W \rangle$).
Hence, γ is in all extensions of $\Delta' := \langle D, W \cup \{\alpha\} \rangle$.
The definition of extensions is invariant under replacing any formula by an equivalent formula.
Thus, $\langle D, W \cup \{\beta\} \rangle$ has exactly the same extensions as Δ' , and γ is in every one of them. Hence, $\beta \sim_{\Delta} \gamma$. \square

Desirable properties: Right Weakening

Right Weakening (RW):

$$\frac{\models \alpha \rightarrow \beta, \gamma \vdash \alpha}{\gamma \vdash \beta}$$

- **Rationale:** If something can be concluded normally, then everything classically implied should also be concluded normally.
- **Example:** Assume that Mary goes **normally implies** Clive goes **and** John goes. Then we would expect that Mary goes **normally implies** Clive goes.
- From (Ref) & (RW) **Supraclassicality** follows:

$$\alpha \vdash \alpha + \frac{\models \alpha \rightarrow \beta, \alpha \vdash \alpha}{\alpha \vdash \beta} \Rightarrow \frac{\alpha \models \beta}{\alpha \vdash \beta}$$

Right Weakening in default logic

Proposition

Default logic satisfies Right Weakening.

Proof.

Assume $\models \alpha \rightarrow \beta$ and $\gamma \vdash_{\Delta} \alpha$ (with $\Delta = \langle D, W \rangle$).
Hence, α is in each extension E of the default theory $\langle D, W \cup \{\gamma\} \rangle$.
Since extensions are closed under logical consequence, β must also be in each extension of $\langle D, W \cup \{\gamma\} \rangle$.
Hence, $\gamma \vdash_{\Delta} \beta$ □

Desirable properties: Cut

Cut:

$$\frac{\alpha \vdash \beta, \alpha \wedge \beta \vdash \gamma}{\alpha \vdash \gamma}$$

- **Rationale:** If part of the premise is plausibly implied by another part of the premise, then the latter is enough for the plausible conclusion.
- **Example:** Assume that John goes **normally implies** Mary goes. Assume further that John goes **and** Mary goes **normally implies** Clive goes. Then we would expect that John goes **normally implies** Clive goes.

Cut in default logic

Proposition

Default logic satisfies Cut.

Proof idea.

Assume $\alpha \vdash_{\Delta} \beta$ (with $\Delta = \langle D, W \rangle$). Hence β is contained in each extension of $\Delta' := \langle D, W \cup \{\alpha\} \rangle$. Show that every extension E of Δ' is also an extension of $\Delta'' = \langle D, W \cup \{\alpha \wedge \beta\} \rangle$.

- Consistency of justifications of defaults is tested against E both in the $W \cup \{\alpha\}$ case and in the $W \cup \{\alpha \wedge \beta\}$ case.
- The preconditions that are derivable when starting from $W \cup \{\alpha\}$ are also derivable when starting from $W \cup \{\alpha \wedge \beta\}$.
- $W \cup \{\alpha \wedge \beta\}$ does not allow for deriving further preconditions because also in the $W \cup \{\alpha\}$ case at some point β is derived.

Hence, because γ belongs to all extensions of Δ'' ($\alpha \wedge \beta \vdash \gamma$), it also belongs to all extensions of Δ' ($\alpha \vdash \gamma$). □

Desirable properties: Cautious Monotonicity



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Cautious Monotonicity (CM):

$$\frac{\alpha \vdash \beta, \alpha \vdash \gamma}{\alpha \wedge \beta \vdash \gamma}$$

- **Rationale:** In general, adding new premises may cancel some conclusions.
However, existing conclusions may be added to the premises without canceling any conclusions!
- **Example:** Assume that
Mary goes **normally implies** Clive goes and
Mary goes **normally implies** John goes.
Mary goes **and** Jack goes might not **normally imply** that John goes.
However, Mary goes and Clive goes should **normally imply** that John goes.

Cautious Monotonicity in default logic



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Proposition

Default logic *does not* satisfy Cautious Monotonicity.

Proof.

Consider the default theory $\langle D, W \rangle$ with

$$D = \left\{ \frac{a : g}{g}, \frac{g : b}{b}, \frac{b : \neg g}{\neg g} \right\} \text{ and } W = \{a\}.$$

$E = \text{Th}(\{a, b, g\})$ is the only extension of $\langle D, W \rangle$ and thus both b and g follow skeptically (i.e., we have $a \vdash_{\langle D, W \rangle} b$ and $a \vdash_{\langle D, W \rangle} g$).

For $\langle D, \{a \wedge b\} \rangle$ also $\text{Th}(\{a, b, \neg g\})$ is an extension, and thus g does not follow skeptically (i.e., $a \wedge b \not\vdash_{\langle D, \{a \wedge b\} \rangle} g$). \square

Cumulativity



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Lemma

Rules (Cut) & (CM) can be equivalently stated as follows:

If $\alpha \vdash \beta$, then the sets of plausible conclusions from α and $\alpha \wedge \beta$ are identical.

This property is called **Cumulativity**.

Proof.

\Rightarrow : Assume that we may apply both rules (Cut) and (CM) and assume $\alpha \vdash \beta$.

Assume further that $\alpha \vdash \gamma$. By applying (CM), we obtain $\alpha \wedge \beta \vdash \gamma$. Similarly, by applying (Cut), from $\alpha \wedge \beta \vdash \gamma$ it follows $\alpha \vdash \gamma$.

Hence the plausible conclusions from α and $\alpha \wedge \beta$ are the same.

\Leftarrow : Assume **Cumulativity** and $\alpha \vdash \beta$. Now we can derive both rules (Cut) and (CM). \square

The System C



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

1 Reflexivity

$$\overline{\alpha \vdash \alpha}$$

2 Left Logical Equivalence

$$\frac{\models \alpha \leftrightarrow \beta, \alpha \vdash \gamma}{\beta \vdash \gamma}$$

3 Right Weakening

$$\frac{\models \alpha \rightarrow \beta, \gamma \vdash \alpha}{\gamma \vdash \beta}$$

4 Cut

$$\frac{\alpha \vdash \beta, \alpha \wedge \beta \vdash \gamma}{\alpha \vdash \gamma}$$

5 Cautious Monotonicity

$$\frac{\alpha \vdash \beta, \alpha \vdash \gamma}{\alpha \wedge \beta \vdash \gamma}$$

Derived rules in C

■ Equivalence:

$$\frac{\alpha \vdash \beta, \beta \vdash \alpha, \alpha \vdash \gamma}{\beta \vdash \gamma}$$

■ And:

$$\frac{\alpha \vdash \beta, \alpha \vdash \gamma}{\alpha \vdash \beta \wedge \gamma}$$

■ MPC:

$$\frac{\alpha \vdash \beta \rightarrow \gamma, \alpha \vdash \beta}{\alpha \vdash \gamma}$$

Introduction
Motivation
Properties
Derived Rules in C
Undesirable
Properties
Reasoning
Semantics
Preferential
Reasoning
Literature

Derived rules: proofs

Proof (Equivalence).

Assumption: $\alpha \vdash \beta, \beta \vdash \alpha, \alpha \vdash \gamma$
Cautious Monotonicity: $\alpha \wedge \beta \vdash \gamma$
Left L Equivalence: $\beta \wedge \alpha \vdash \gamma$
Cut: $\beta \vdash \gamma$ □

Proof (And).

Assumption: $\alpha \vdash \beta, \alpha \vdash \gamma$
Cautious Monotonicity: $\alpha \wedge \beta \vdash \gamma$
propositional logic: $\alpha \wedge \beta \wedge \gamma \models \beta \wedge \gamma$
Supraclassicality: $\alpha \wedge \beta \wedge \gamma \vdash \beta \wedge \gamma$
Cut: $\alpha \wedge \beta \vdash \beta \wedge \gamma$
Cut: $\alpha \vdash \beta \wedge \gamma$ □

MPC is an exercise.

Introduction
Motivation
Properties
Derived Rules in C
Undesirable
Properties
Reasoning
Semantics
Preferential
Reasoning
Literature

Undesirable properties: Monotonicity and Contraposition

■ Monotonicity: $\frac{\models \alpha \rightarrow \beta, \beta \vdash \gamma}{\alpha \vdash \gamma}$

- *Example:* Let us assume that
John goes **normally implies** Mary goes.
Now we will probably not expect that
John goes **and** Joan (who is not in talking terms with
Mary) goes **normally implies** Mary goes.


■ Contraposition: $\frac{\alpha \vdash \beta}{\neg \beta \vdash \neg \alpha}$

- *Example:* Let us assume that
John goes **normally implies** Mary goes.
Would we expect that
Mary does not go **normally implies** John does not go?
What if John goes always?

Introduction
Motivation
Properties
Derived Rules in C
Undesirable
Properties
Reasoning
Semantics
Preferential
Reasoning
Literature

Undesirable properties: Monotonicity


$\alpha \models \beta, \beta \vdash \gamma$, but not $\alpha \vdash \gamma$ — pictorially:



Introduction
Motivation
Properties
Derived Rules in C
Undesirable
Properties
Reasoning
Semantics
Preferential
Reasoning
Literature

Undesirable properties: Contraposition

$\alpha \sim \beta$, but not $\neg\beta \sim \neg\alpha$ — pictorially:



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Undesirable properties: Transitivity & EHD

■ Transitivity:

$$\frac{\alpha \sim \beta, \beta \sim \gamma}{\alpha \sim \gamma}$$

- **Example:** Let us assume that
John goes **normally implies** Mary goes and
Mary goes **normally implies** Jack goes.
Now, should John goes **normally imply** that Jack goes?
What, if John goes very seldom?


■ Easy Half of the Deduction Theorem (EHD):

$$\frac{\alpha \sim \beta \rightarrow \gamma}{\alpha \wedge \beta \sim \gamma}$$

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Undesirable properties: Transitivity


$\alpha \sim \beta, \beta \sim \gamma$, but not $\alpha \sim \gamma$ — pictorially:



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Undesirable properties: EHD

$\alpha \sim \beta \rightarrow \gamma$, but not $\alpha \wedge \beta \sim \gamma$ — pictorially:



Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Monotonicity vs EHD



Theorem

In the presence of the rules in system **C**, the rules *Monotonicity* and *EHD* are equivalent.

Proof.

Monotonicity \Rightarrow EHD:

- $\alpha \sim \beta \rightarrow \gamma$ (assumption)
- $\alpha \wedge \beta \sim \beta \rightarrow \gamma$ (Monotonicity)
- $\alpha \wedge \beta \sim \alpha \wedge \beta$ (Ref)
- $\alpha \wedge \beta \sim \beta$ (RW)
- $\alpha \wedge \beta \sim \gamma$ (MPC)

Monotonicity \Leftarrow EHD:

- $\models \alpha \rightarrow \beta, \beta \sim \gamma$ (assumption)
- $\beta \sim \alpha \rightarrow \gamma$ (RW)
- $\beta \wedge \alpha \sim \gamma$ (EHD)
- $\alpha \sim \gamma$ (LLE)

□

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Monotonicity vs Transitivity



Theorem

In the presence of the rules in system **C**, the rules *Monotonicity* and *Transitivity* are equivalent.

Proof.

Monotonicity \Rightarrow Transitivity:

- $\alpha \sim \beta, \beta \sim \gamma$ (assumption)
- $\alpha \wedge \beta \sim \gamma$ (Monotonicity)
- $\alpha \sim \gamma$ (Cut)

Monotonicity \Leftarrow Transitivity:

- $\models \alpha \rightarrow \beta, \beta \sim \gamma$ (assumption)
- $\alpha \models \beta$ (deduction theorem)
- $\alpha \sim \beta$ (Supraclassicality)
- $\alpha \sim \gamma$ (Transitivity)

□

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

Contraposition?



Theorem

In the presence of *Right Weakening*, *Contraposition* implies *Monotonicity*.

Proof.

- $\models \alpha \rightarrow \beta, \beta \sim \gamma$ (assumption)
- $\neg \gamma \sim \neg \beta$ (Contraposition)
- $\models \neg \beta \rightarrow \neg \alpha$ (classical contraposition)
- $\neg \gamma \sim \neg \alpha$ (RW)
- $\alpha \sim \gamma$ (Contraposition)

□

Note: *Monotonicity* does not imply *Contraposition*, even in the presence of all rules of system **C**!

Introduction
Motivation
Properties
Derived Rules in C
Undesirable Properties
Reasoning
Semantics
Preferential Reasoning
Literature

2 Reasoning



Introduction
Reasoning
Semantics
Preferential Reasoning
Literature

- How do we **reason** with \sim from φ to ψ ?
- **Assumption**: We have some (finite) set K of **conditional statements** of the form $\alpha \sim \beta$.
The question is: Assuming the statements in K , is it plausible to conclude ψ given φ ?
- **Idea**: We consider **all** cumulative consequence relations that contain K .
Cumulative consequence relation: any relation \sim between propositional logic formulae that is closed under the rules of system **C**.
- **Remark**: It suffices to consider only the **minimal** cumulative consequence relation containing K ...

Introduction
Reasoning
Semantics
Preferential Reasoning
Literature

Lemma

The set of cumulative consequence relations is closed under (arbitrary) intersections.

Proof.

Let \sim_1 and \sim_2 be cumulative consequence relations. We have to show that $\sim_1 \cap \sim_2$ is a cumulative consequence relation, that is, it is closed under all the rules of system **C**.
Take any instance of any of the rules. If the preconditions are satisfied by \sim_1 and \sim_2 , then the consequence is trivially also satisfied by both. A similar argument works if we consider an arbitrary family of consequence relations. \square

Introduction
Reasoning
Semantics
Preferential Reasoning
Literature

Theorem

For each set of conditional statements K , there exists a unique minimal cumulative consequence relation containing K .

Proof.

From the previous lemma it is clear that the intersection of all the cumulative consequence relations containing K is already such a cumulative consequence relation.
Obviously, there cannot be two distinct such minimal relations. \square

This relation is called the **cumulative closure** K^C of K .

Introduction
Reasoning
Semantics
Preferential Reasoning
Literature

- Cumulative Models
- Consequence Relations

Introduction
Reasoning
Semantics
Cumulative Models
Consequence Relations
Preferential Reasoning
Literature

Cumulative models – informally



- We will now try to characterize cumulative reasoning model-theoretically.
- **Idea:** Cumulative models consist of states ordered by a preference relation.
- States characterize beliefs.
- The preference relation, \prec , expresses the normality of the beliefs.
We read $s \prec t$ as: state s is preferred to/more normal than state t .
- We say: $\alpha \vdash \beta$ is **accepted** in a model if in all most preferred states in which α is true also β is true.

Introduction
Reasoning
Semantics
Cumulative Models
Consequence Relations
Preferential Reasoning
Literature

Preference relation



We consider an arbitrary binary relation \prec on a given set of states S .

Later, we will assume that \prec has particular properties, e.g., that \prec is irreflexive, asymmetric, transitive, a partial order, ...
... but currently we make no such restrictions.

We need a condition on state sets claiming that each state is, or is related to, a most preferred state.

Definition (Smoothness)

Let $P \subseteq S$.

- We say that $s \in P$ is **minimal in P** if $s' \not\prec s$ for each $s' \in P$.
- P is called **smooth** if for each $s \in P$, either s is minimal in P or there exists an s' such that s' is minimal in P and $s' \prec s$.

Introduction
Reasoning
Semantics
Cumulative Models
Consequence Relations
Preferential Reasoning
Literature

Cumulative Models – formally



Let \mathcal{U} be the set of all possible worlds (i.e., propositional interpretations).

- A **cumulative model** is a triple $W = \langle S, I, \prec \rangle$ such that
 - 1 S is a set of states,
 - 2 I is a mapping $I : S \rightarrow 2^{\mathcal{U}}$, and
 - 3 \prec is an arbitrary binary relation on S
 such that the **smoothness condition** is satisfied (see below).
- A state $s \in S$ **satisfies** a formula α ($s \models \alpha$) if $m \models \alpha$ for each propositional interpretation $m \in I(s)$.
The set of states satisfying α is denoted by $\hat{\alpha}$.
- **Smoothness condition:** A cumulative model satisfies this condition if for all formulae α , $\hat{\alpha}$ is **smooth**.

Introduction
Reasoning
Semantics
Cumulative Models
Consequence Relations
Preferential Reasoning
Literature

Consequence relation induced by a cumulative model



A cumulative model W induces a consequence relation \vdash_W as follows:

$$\alpha \vdash_W \beta \text{ iff } s \models \beta \text{ for every minimal } s \text{ in } \hat{\alpha}.$$

Example

Model $W = \langle \{s_1, s_2, s_3\}, I, \prec \rangle$ with $s_1 \prec s_2, s_2 \prec s_3, s_1 \prec s_3$

$$I(s_1) = \{\{\neg p, b, f\}\}$$

$$I(s_2) = \{\{p, b, \neg f\}\}$$

$$I(s_3) = \{\{\neg p, \neg b, f\}, \{\neg p, \neg b, \neg f\}\}$$

Does W satisfy the smoothness condition?

$\neg p \wedge \neg b \vdash f$? N Also: $\neg p \wedge \neg b \not\vdash \neg f$!

$p \vdash \neg f$? Y

$\neg p \vdash f$? Y

Introduction
Reasoning
Semantics
Cumulative Models
Consequence Relations
Preferential Reasoning
Literature

Soundness 1

Theorem

If W is a cumulative model, then \vdash_W is a cumulative consequence relation.

Proof.

- **Reflexivity:** satisfied.
- **LLE:** satisfied.
- **RW:** satisfied.
- **Cut:** $\alpha \vdash_W \beta, \alpha \wedge \beta \vdash_W \gamma \Rightarrow \alpha \vdash_W \gamma$. Assume that all minimal elements of $\hat{\alpha}$ satisfy β , and all minimal elements of $\widehat{\alpha \wedge \beta}$ satisfy γ . Every minimal element of $\hat{\alpha}$ satisfies $\alpha \wedge \beta$. Since $\widehat{\alpha \wedge \beta} \subseteq \hat{\alpha}$, all minimal elements of $\hat{\alpha}$ are also minimal elements of $\widehat{\alpha \wedge \beta}$. Hence $\alpha \vdash_W \gamma$.

Soundness 2

Proof continues...

- **Cautious Monotonicity:** $(\alpha \vdash \beta, \alpha \vdash \gamma \Rightarrow \alpha \wedge \beta \vdash \gamma)$

Assume $\alpha \vdash_W \beta$ and $\alpha \vdash_W \gamma$. We have to show: $\alpha \wedge \beta \vdash_W \gamma$, i.e., $s \models \gamma$ for all minimal $s \in \widehat{\alpha \wedge \beta}$.

Clearly, every minimal $s \in \widehat{\alpha \wedge \beta}$ is in $\hat{\alpha}$.

We show that every minimal $s \in \widehat{\alpha \wedge \beta}$ is **minimal** in $\hat{\alpha}$.

Assumption: There is s that is minimal in $\widehat{\alpha \wedge \beta}$, but not minimal in $\hat{\alpha}$. Because of **smoothness** there is minimal $s' \in \hat{\alpha}$ such that $s' \prec s$. We know, however, that $s' \models \beta$, which means that $s' \in \widehat{\alpha \wedge \beta}$. Hence s is not minimal in $\widehat{\alpha \wedge \beta}$. **Contradiction!**

Hence s must be minimal in $\hat{\alpha}$, and therefore $s \models \gamma$. Because this is true for all minimal elements in $\widehat{\alpha \wedge \beta}$, we get $\alpha \wedge \beta \vdash_W \gamma$. \square

Consequence: counterexamples

Now we have a **method** for showing that a principle does not hold for cumulative consequence relations:

... construct a **cumulative model** that falsifies the principle.

Contraposition: $\alpha \vdash \beta \Rightarrow \neg\beta \vdash \neg\alpha$

$$W = \langle S, I, \prec \rangle$$

$$S = \{s_1, s_2\}$$

$$s_i \not\prec s_j \quad \forall s_i, s_j \in S$$

$$I(s_1) = \{\{a, b\}\}$$

$$I(s_2) = \{\{a, \neg b\}, \{\neg a, \neg b\}\}$$

W is a cumulative model with $a \vdash_W b$, but $\neg b \not\vdash_W \neg a$.

Completeness?

- Each cumulative model W **induces** a cumulative consequence relation \vdash_W .
- **Problem:** Can we generate all cumulative consequence relations in this way?
- We can! There is a **representation theorem**:

Theorem (Representation of cumulative consequence)

A consequence relation is cumulative if and only if it is induced by some cumulative model.

\rightsquigarrow Cumulative consequence can be characterized independently from the set of inference rules.

Transitivity of the preference relation?

- Could we strengthen the preference relation to **transitive** relations without sacrificing anything?
No!
- In such models, the following additional principle called **Loop** is valid:

$$\frac{\alpha_0 \sim \alpha_1, \alpha_1 \sim \alpha_2, \dots, \alpha_k \sim \alpha_0}{\alpha_0 \sim \alpha_k}$$

- For the system **CL** = **C** + (**Loop**) and cumulative models with transitive preference relations, we could prove another **representation theorem**.

The Or Rule

Or rule:

$$\frac{\alpha \sim \gamma, \beta \sim \gamma}{\alpha \vee \beta \sim \gamma}$$

Not valid in system **C**. **Counterexample**:

$$W = \langle S, I, \prec \rangle$$

$$S = \{s_1, s_2, s_3\}, s_i \not\prec s_j \forall s_i, s_j \in S$$

$$I(s_1) = \{\{a, b, c\}, \{a, \neg b, c\}\}$$

$$I(s_2) = \{\{a, b, c\}, \{\neg a, b, c\}\}$$

$$I(s_3) = \{\{a, b, \neg c\}, \{a, \neg b, \neg c\}, \{\neg a, b, \neg c\}\}$$

$a \sim_W c, b \sim_W c$, but not $a \vee b \sim_W c$.

Note: Or is not valid in default logic.

4 Preferential Reasoning

- Preferential Relations

System P

- System **P** contains all rules of **C** and the Or rule.
- A consequence relation that satisfies **P** is called **preferential**.
- Derived rules in **P**:
 - Hard half of the deduction theorem (**S**):

$$\frac{\alpha \wedge \beta \sim \gamma}{\alpha \sim \beta \rightarrow \gamma}$$

- Proof by case analysis (**D**):

$$\frac{\alpha \wedge \neg \beta \sim \gamma, \alpha \wedge \beta \sim \gamma}{\alpha \sim \gamma}$$

- **D** and Or are equivalent in the presence of the rules in **C**.

Definition

A cumulative model $W = \langle S, I, \prec \rangle$ such that \prec is a **strict partial order** (irreflexive and transitive) and $|I(s)| = 1$ for all $s \in S$ is called a **preferential model**.

Theorem (Soundness)

The consequence relation \vdash_W induced by a preferential model is preferential.

Proof.

Since W is cumulative, we only have to verify that **Or** holds. Note that in preferential models we have $\widehat{\alpha \vee \beta} = \widehat{\alpha} \cup \widehat{\beta}$. Suppose $\alpha \vdash_W \gamma$ and $\beta \vdash_W \gamma$. Because of the above equation, each minimal state of $\widehat{\alpha \vee \beta}$ is minimal in $\widehat{\alpha} \cup \widehat{\beta}$. Since γ is satisfied in all minimal states in $\widehat{\alpha} \cup \widehat{\beta}$, γ is also satisfied in all minimal states of $\widehat{\alpha \vee \beta}$. Hence $\alpha \vee \beta \vdash_W \gamma$. \square

Theorem (Representation of preferential consequence)

A consequence relation is preferential if and only if it is induced by a preferential model.

Proof.

Similar to the one for **C**. \square

System

C
 Reflexivity
 Left Logical Equivalence
 Right Weakening
 Cut
 Cautious Monotonicity

CL

+ Loop

P

+ Or

Models

States: sets of worlds
 Preference relation: arbitrary
 Models must be smooth

Preference relation: strict partial order

States: singletons

Strengthening the consequence relation

- System **C** and System **P** do not produce many of the inferences one would hope for:

Given $K = \{Bird \sim Flies\}$ one cannot conclude $Red \wedge Bird \sim Flies!$

- In general, adding information that is **irrelevant** cancels the plausible conclusions.
 \Rightarrow Cumulative and Preferential consequence relations are **too nonmonotonic**.
- The plausible conclusions have to be strengthened!

Introduction
Reasoning
Semantics
Preferential Reasoning
Preferential Relations
Literature

Strengthening the consequence relations

- The rules so far seem to be reasonable: one cannot think of rules of the same form (if we have some plausible implications, other plausible implications should hold) that could be added.
- However, there are other types of rules one might want add.
- **Disjunctive Rationality**:

$$\frac{\alpha \not\sim \gamma, \beta \not\sim \gamma}{\alpha \vee \beta \not\sim \gamma}$$

- **Rational Monotonicity**:

$$\frac{\alpha \sim \gamma, \alpha \not\sim \neg\beta}{\alpha \wedge \beta \sim \gamma}$$

- **Note**: Consequence relations obeying these rules are not closed under intersection, which is a problem.

Introduction
Reasoning
Semantics
Preferential Reasoning
Preferential Relations
Literature

Summary

- Instead of **ad hoc** extensions of the logical machinery, analyze the properties of nonmonotonic consequence relations.
- Correspondence between rule system and models for System **C**, and for System **P** could also be established wrt. a probabilistic semantics.
- Irrelevant information poses a problem. Solution approaches: rational monotonicity, maximum entropy approach.

Introduction
Reasoning
Semantics
Preferential Reasoning
Preferential Relations
Literature

5 Literature

Introduction
Reasoning
Semantics
Preferential Reasoning
Literature



Sarit Kraus, Daniel Lehmann, and Menachem Magidor.

Nonmonotonic reasoning, preferential models and cumulative logics.

[Artificial Intelligence](#), 44:167–207, 1990.

Introduces cumulative consequence relations.



Daniel Lehman and Menachem Magidor.

What does a conditional knowledge base entail?

[Artificial Intelligence](#), 55:1–60, 1992.

Introduces rational consequence relations.



Dov M. Gabbay.

Theoretical foundations for non-monotonic reasoning in expert systems.

In K. R. Apt, editor, **Proceedings NATO Advanced Study Institute on Logics and Models of Concurrent Systems**, pages 439–457.

Springer-Verlag, Berlin, Heidelberg, New York, 1985.

First to consider abstract properties of nonmonotonic consequence relations.

Introduction

Reasoning

Semantics

Preferential
Reasoning

Literature



Judea Pearl.

Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference,

Morgan Kaufmann Publishers, 1988.

One section on ϵ -semantics and maximum entropy.



Yoav Shoham.

Reasoning about Change.

MIT Press, Cambridge, MA, 1988.

Introduces the idea of preferential models.

Introduction

Reasoning

Semantics

Preferential
Reasoning

Literature