

Principles of Knowledge Representation and Reasoning

Dynamics of belief

Albert-Ludwigs-Universität Freiburg



Bernhard Nebel, Stefan Wölfl, and Julien Hué

Winter Semester 2012/2013

1 Introduction

- Link between revision and update

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Propositional logic flaws:

- The world is not always static.
- The knowledge about the world is sometimes uncertain or imprecise

Therefore:

- Need the possibility to incorporate new (possibly contradictory) beliefs;
- Need to take into account change in the world;

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Plato - Theaetetus: A knowledge (a rightful opinion) is a piece of

1 Justified True Belief

Agrippa's trilemma - A problem with the justification:

- 1 Either the justification stops to some unjustified belief;
- 2 The justification is infinite (Socrates' clouds);
- 3 The justification is supported by affirmations it is supposed to justify (Baron Münchhausen's hair).

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Three solutions:

Foundationalism Allow for unjustified beliefs

- Formalization issues
- Humans don't keep track of sources
- **TMS System**

“Infinetism” Allow for infinite justification

- Does it really make sense?

Coherentism Allow for circular justifications

- What is a solid belief?
- **Belief revision/update**

- In any cases, information is extremely important and should not be discarded carelessly.

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Arrow's impossibility theorem - there is no voting system which respects:

- Non-dictatorship
(all voters should be taken into account);
- Universality
(complete and deterministic ranking);
- Independance of irrelevant alternatives
(ranking between x and y depends only on x and y);
- Pareto efficiency
(if all preferences states $x < y$, then so must the results).

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

- We have a theory about the world, and the new information is meant to **correct** our theory
- ⇒ **belief revision**: change your belief state minimally in order to accommodate the new information
- We have a (supposedly) correct theory about the current state of the world, and the new information is meant to record a **change** in the world
- ⇒ **belief update**: incorporate the change by assuming that the world has changed minimally

Assume the new information is consistent with our old beliefs.

- In case of **belief revision**, we would like to add the new information monotonically to our old beliefs.
- For **belief update** this is not necessarily the case.
 - Assume we know that the **door is open or the window is open**.
 - Assume we learn that the world has changed and the **door is now closed**.
- In this case, we do not want to add this information monotonically to our theory, since we would be forced to conclude that **the window is open**.

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

What are the criteria for definition of a belief revision operation?

Gärdenfors and Rott - belief revision (1995):

- 1 How are beliefs represented?
- 2 What is the relation between beliefs represented explicitly in the belief base and beliefs which can be derived from them?
- 3 In the face of a contradiction, how to deal with both new and old information?

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Belief base, belief set or interpretation?

General assumption:

- A **belief set** is a deductively closed theory, i.e., $K = \text{Cn}(K)$ with Cn the **consequence operator**
- \mathcal{L} : logical language (propositional logic)
- $\text{Th}_{\mathcal{L}}$: set of deductively closed theories (or belief sets) over \mathcal{L}

Belief change operations

Monotonic addition: $+: \text{Th}_{\mathcal{L}} \times \mathcal{L} \rightarrow \text{Th}_{\mathcal{L}}$
 $K + \psi = \text{Cn}(K \cup \{\psi\})$

Revision: $\dot{+}: \text{Th}_{\mathcal{L}} \times \mathcal{L} \rightarrow \text{Th}_{\mathcal{L}}$

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

Consider $K = \{a, b\}$ and $K' = \{a \wedge b\}$. What is happening to $K \dot{+} \{\neg a\}$?

Semantic

- No difference between K and K'

a	b	\mathcal{I}
0	0	0
0	1	0
1	0	0
1	1	1

Syntactic

- $X = \{b\}$ is the only maximal subset of K s.t. $X \cup \{\neg a\}$ is consistent.
- $X' = \emptyset$ is the only maximal subset of K' s.t. $X' \cup \{\neg a\}$ is consistent.

Introduction

Link between
revision and update

Belief
revision

Several
sources -
belief
merging

Bibliography

2 Belief revision

- Formal properties
- Standard revision operations
- Semantic approaches

Introduction

**Belief
revision**

Formal properties

Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

What is a good revision operator?

- Consistency: a revision has to produce a consistent set of beliefs;
- Minimality of change: a revision has to change the fewest possible beliefs;
- Priority to the new information: the 'new' information is considered more important than the 'old' one.

Introduction

Belief
revision

Formal properties

Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

AGM postulates:

- ($\dot{+}$ 1) $K \dot{+} \varphi \in \text{Th}_{\mathcal{L}}$;
- ($\dot{+}$ 2) $\varphi \in K \dot{+} \varphi$;
- ($\dot{+}$ 3) $K \dot{+} \varphi \subseteq K + \varphi$;
- ($\dot{+}$ 4) If $\neg\varphi \notin K$, then $K + \varphi \subseteq K \dot{+} \varphi$;
- ($\dot{+}$ 5) $K \dot{+} \varphi = \text{Cn}(\perp)$ only if $\vdash \neg\varphi$;
- ($\dot{+}$ 6) If $\vdash \varphi \leftrightarrow \psi$ then $K \dot{+} \varphi = K \dot{+} \psi$;
- ($\dot{+}$ 7) $K \dot{+} (\varphi \wedge \psi) \subseteq (K \dot{+} \varphi) + \psi$;
- ($\dot{+}$ 8) If $\neg\psi \notin K \dot{+} \varphi$,
then $(K \dot{+} \varphi) + \psi \subseteq K \dot{+} (\varphi \wedge \psi)$.

Introduction

Belief
revision

Formal properties

Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

Revision can be defined in terms of two suboperations.

- $+$ (**expansion**) denotes the simple union of beliefs;
- $-$ (**contraction**) denotes the removal of information contradicting the input.

The Levi identity

$$K \dot{+} \varphi \equiv Cn[(K - \neg\varphi) + \varphi]$$

Example

$$K = \{a, a \rightarrow b\} \quad \varphi = \{\neg b\}?$$

$$K - \neg\varphi = \{a\} \text{ or } \{a \rightarrow b\}$$

$$K \dot{+} \neg\varphi = \{a, \neg b\} \text{ or } \{a \rightarrow b, \neg b\}$$

Introduction

Belief
revision

Formal properties

Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

Definition

We denote by $K \perp \varphi$ the set of maximal (wrt set-theoretic inclusion) subsets J of K such that $J \not\models \varphi$.

Definition

Full-meet contraction is defined by $K - \varphi = \bigcap (K \perp \varphi)$.

Is full-meet contraction reasonable?

- ▶ No! It is far too cautious.
- ▶ It can nevertheless be used as a lower bound to any reasonable operator.

$K \dot{+} \varphi = \bigcap (K \perp \varphi) + \varphi$ is referred to as the **full-meet revision**.

Introduction

Belief
revision

Formal properties
Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

Proposition

Full-meet revision respects all AGM postulates.

Proof

($\dot{+}$ 1) and ($\dot{+}$ 2) are true by construction

($\dot{+}$ 3) Two cases: (1) If $K + \varphi$ is consistent then $K - \varphi = K$ and $K \dot{+} \varphi = K + \varphi$. (2) If $K + \varphi$ is inconsistent then $K + \varphi = \text{Cn}(\perp)$ and $K \dot{+} \varphi \subseteq K + \varphi$.

($\dot{+}$ 4) Because $K \not\vdash \neg\varphi$ then $K \perp \varphi = \{K\}$ and thus $K \dot{+} \varphi = K + \varphi$.

($\dot{+}$ 5) $K \dot{+} \varphi = \text{Cn}(\bigcap_{\alpha \in (K \perp \varphi)} \alpha \cup \varphi)$. But $\forall \alpha, \alpha \cup \varphi \not\vdash \perp$, therefore $\bigcap_{\alpha \in (K \perp \varphi)} \alpha \cup \varphi \not\vdash \perp$ (as PL is monotonic).

($\dot{+}$ 6) Lets assume that $\alpha \in K \perp \varphi$ but $\alpha \notin K \perp \psi$. Two cases: (1) $\alpha \cup \psi \vdash \perp \xrightarrow{(\varphi \leftrightarrow \psi)} \alpha \cup \varphi \vdash \perp$ which is not possible. (2) $\exists \beta$ s.t. $\alpha \subsetneq \beta$ and $\beta \cup \psi \not\vdash \perp \xrightarrow{(\varphi \leftrightarrow \psi)} \beta \cup \varphi \not\vdash \perp$ which is not possible.

($\dot{+}$ 7) and ($\dot{+}$ 8) Left as exercises...

Introduction

Belief
revision

Formal properties
Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

On the other side, one can ask for the principle of minimality to be strictly respected.

Definition

A selection function for K is a function γ such that for all sentences φ :

- 1 If $K \perp \varphi$ is non-empty, then $\gamma(K \perp \varphi)$ is a non-empty subset of $K \perp \varphi$, and
- 2 If $K \perp \varphi$ is empty, then $\gamma(K \perp \varphi) = \{K\}$.

Definition

Maxichoice contraction is defined as $K - \varphi = \gamma(K \perp \varphi)$ where γ is a selection function.

Introduction

Belief
revision

Formal properties
Standard revision
operations
Semantic
approaches

Several
sources -
belief
merging

Bibliography

Maxi-choice can be too bold: there is sometimes no reason to trust one piece more than one another.

Definition

A partial-meet revision operation is an operation defined as:

$$K \dot{+} \varphi = \bigcap \gamma(K \perp \varphi) + \varphi$$

Seems to be a good compromise between full-meet and maxi-choice

Introduction

Belief
revision

Formal properties
Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

Definition

The Dalal revision operation, denoted by $\dot{+}_D$, is defined as:

$$K \dot{+}_D \varphi = \min(\text{extMod}(\varphi), \leq_K)$$

where d_H is the Hamming Distance and

$$\alpha \leq_K \beta \text{ iff } \exists \omega \in \text{extMod}(K), \forall \omega' \in \text{extMod}(K), d_H(\alpha, \omega) \leq d_H(\beta, \omega')$$

Example

	<i>a</i>	<i>b</i>	<i>c</i>
\mathcal{I}_{φ_1}	0	0	0
\mathcal{I}_{φ_2}	0	0	1
	0	1	0
\mathcal{I}_{K_1}	0	1	1
	1	0	0
\mathcal{I}_{K_2}	1	0	1
	1	1	0
\mathcal{I}_{K_3}	1	1	1

Let $\varphi = \{\neg a, \neg b\}$ and $K = \{(a \vee b) \wedge c\}$:

$$d(\mathcal{I}_{\varphi_1}, \mathcal{I}_{K_1}) = 2 \quad d(\mathcal{I}_{\varphi_2}, \mathcal{I}_{K_1}) = 1$$

$$d(\mathcal{I}_{\varphi_1}, \mathcal{I}_{K_2}) = 2 \quad d(\mathcal{I}_{\varphi_2}, \mathcal{I}_{K_2}) = 1$$

$$d(\mathcal{I}_{\varphi_1}, \mathcal{I}_{K_3}) = 3 \quad d(\mathcal{I}_{\varphi_2}, \mathcal{I}_{K_3}) = 2$$

Introduction

Belief
revision

Formal properties
Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

Formula-based approaches

The question does Ψ belongs to $K \dot{+} \varphi$ (if $\dot{+}$ is a full-meet revision operator) is $\Delta_2^P - (\Sigma_1^P \cup \Pi_1^P)$ provided that $\text{NP} \neq \text{co-NP}$.

proof

If $\dot{+}$ is a full-meet revision, $\Psi \in \text{Cn}(K) \dot{+} \varphi$ can be solved by the following algorithm: if $K \not\models \neg\Psi$, then $K \cup \Psi \models \varphi$ else $\Psi \models \varphi \rightarrow$ Membership in Δ_2^P .

Furthermore, SAT can be polynomially transformed to full-meet revision by solving $\Psi \in \text{Cn}(\Psi) \dot{+} \top$ and UNSAT can be polynomially transform to full-meet revision by solving $\perp \in \text{Cn}(\emptyset) \dot{+} \Psi$. Hence, assuming that full-meet revision belongs to both NP and co-NP would lead to $\text{NP} = \text{co-NP}$.

Introduction

Belief
revision

Formal properties
Standard revision
operations

Semantic
approaches

Several
sources -
belief
merging

Bibliography

3 Several sources - belief merging

- Postulational aspects
- Distance-based merging
- Syntactic merging

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

There is not only one source for the information:

- Voting procedure;
- Expert system;
- Distributed databases;
- multisource knowledge acquisition.

Constructing a belief base which represents the several sources and which:

- solves the contradiction;
- reduces the redundancies;
- is consistent.

Introduction

Belief
revision

Several
sources -
belief
merging


Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Merging in the general case



General assumption:

- K_1, \dots, K_n are belief bases;
- $E = \{K_1, \dots, K_n\}$ is a **multi-set** of belief bases and is called a **belief profile**;
- IC is a propositional formula standing for constraints;
- \sqcup stands for multi-set union.

Operation

Belief merging operation: $\Delta : \mathcal{L}^n \times \mathcal{L} \rightarrow \mathcal{L}$

Sometimes also called **fusion** operation.

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

- (KP0) $\Delta_{IC}(E) \models IC$.
- (KP1) If IC is consistent, then $\Delta_{IC}(E)$ is consistent.
- (KP2) If $\bigwedge E \wedge IC$ is consistent, then $\Delta_{IC}(E) = \bigwedge E \wedge IC$.
- (KP3) If $E_1 \equiv E_2$ and $IC_1 \equiv IC_2$, then $\Delta_{IC_1}(E_1) \equiv \Delta_{IC_2}(E_2)$.
- (KP4) If $K_1 \models IC$ and $K_2 \models IC$, then $\Delta_{IC}(K_1 \sqcup K_2) \wedge K_1 \not\models \perp$ implies $\Delta_{IC}(K_1 \sqcup K_2) \wedge K_2 \not\models \perp$.
- (KP5) $\Delta_{IC}(E_1) \wedge \Delta_{IC}(E_2) \models \Delta_{IC}(E_1 \sqcup E_2)$.
- (KP6) If $\Delta_{IC}(E_1) \wedge \Delta_{IC}(E_2)$ is consistent, then $\Delta_{IC}(E_1 \sqcup E_2) \models \Delta_{IC}(E_1) \wedge \Delta_{IC}(E_2)$.
- (KP7) $\Delta_{IC_1}(E) \wedge IC_2 \models \Delta_{IC_1 \wedge IC_2}(E)$.
- (KP8) If $\Delta_{IC_1}(E) \wedge IC_2$ is consistent, then $\Delta_{IC_1 \wedge IC_2}(E) \models \Delta_{IC_1}(E) \wedge IC_2$.

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Arbitration (Arb)

$$\left. \begin{array}{l} \Delta_{IC_1}(K_1) \leftrightarrow \Delta_{IC_2}(K_2) \\ \Delta_{IC_1 \leftrightarrow \neg IC_2}(K_1 \sqcup K_2) \leftrightarrow (IC_1 \leftrightarrow \neg IC_2) \\ IC_1 \neg \vdash IC_2 \\ IC_2 \neg \vdash IC_1 \end{array} \right\} \Rightarrow \Delta_{IC_1 \vee IC_2}(K_1 \sqcup K_2) \leftrightarrow \Delta_{IC_1}(K_1)$$

Majority (Maj)

$$\exists n, \Delta_{IC}(K_1 \sqcup K_2^n) \vdash \Delta_{IC}(K_2)$$

Independence from majority (IM)

$$\forall n, \Delta_{IC}(K_1 \sqcup K_2^n) \leftrightarrow \Delta_{IC}(K_1 \sqcup K_2)$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Theorem

There exists no merging operator satisfying all the KP postulates and (IM).

Proof

Consider $E_1 = \{K, \neg K\}$ and $E_2 = \{K\}$ be two belief profiles.

(IM) leads to $\Delta_{\top}(E_1 \sqcup E_2) = \Delta_{\top}(E_1)$.

(KP4) allows for $\Delta_{\top}(E_1) \not\models K$ and $\Delta_{\top}(E_1) \not\models \neg K$.

From (KP2), we have that $\Delta_{\top}(E_2) \models K$ and thus $\Delta_{\top}(E_1) \wedge \Delta_{\top}(E_2)$ is consistent and from (KP6) we obtain $\Delta_{\top}(E_1 \sqcup E_2) \models \Delta_{\top}(E_1) \wedge \Delta_{\top}(E_2)$, i.e., $\Delta_{\top}(E_1) \models \Delta_{\top}(E_1) \wedge K$ and thus $\Delta_{\top}(E_1) \models K$ contradicting (KP4).

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Theorem

If a merging operator satisfies (KP1) and (KP2) then it can not satisfies (IM) and (Maj) at the same time.

Proof

From (IM) and (Maj), we have for all E_1, K that

$$\Delta_{\top}(E_1 \sqcup K) \leftrightarrow \Delta_{\top}(E_1 \sqcup K^n) \vdash \Delta_{\top}(K).$$

From (KP2), we deduce that $\forall K, \Delta_{\top}(E_1 \sqcup K) \vdash K$.

Consider K' such that $K \wedge K' \vdash \perp$. Then with $E = K'$, we have

$\Delta_{\top}(K' \sqcup K) \vdash K$. And also that $\Delta_{\top}(K \sqcup K') \vdash K'$ and thus that

$\Delta_{\top}(K' \sqcup K) \vdash K \wedge K'$. Finally, $\Delta_{\top}(K' \sqcup K) \vdash \perp$ contradicting (KP1).

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Definition

A syncretic assignment is a function which associates to a belief profile E a pre-order \leq_E over the interpretations such that for every belief profile E, E_1, E_2 and every belief base K, K' the following conditions hold:

- 1 If $\omega \models E$ and $\omega' \models E$ then $\omega \simeq_E \omega'$
- 2 If $\omega \models E$ and $\omega' \not\models E$ then $\omega <_E \omega'$
- 3 If $E_1 \leftrightarrow E_2$ then $\leq_{E_1} = \leq_{E_2}$
- 4 $\forall \omega \models K, \exists \omega' \models K', \omega' \leq_{K \sqcup K'} \omega$
- 5 If $\omega \leq_{E_1} \omega'$ and $\omega \leq_{E_2} \omega'$ then $\omega \leq_{E_1 \sqcup E_2} \omega'$
- 6 If $\omega <_{E_1} \omega'$ and $\omega \leq_{E_2} \omega'$ then $\omega <_{E_1 \sqcup E_2} \omega'$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Definition

A majority syncretic assignment is a syncretic assignment which satisfies the following condition:

$$7 \text{ If } \omega <_{E_2} \omega', \text{ then } \exists n, \omega <_{E_1 \sqcup E_2^n} \omega'$$

Definition

A fair syncretic assignment is a syncretic assignment which satisfies the following condition:

8

$$\left. \begin{array}{l} \omega <_K \omega' \\ \omega <_{K'} \omega'' \\ \omega' \simeq_{K \sqcup K'} \omega'' \end{array} \right\} \Rightarrow \omega <_{K \sqcup K'} \omega'$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography



Theorem

We consider Δ_{IC} a merging operation. Δ_{IC} respects all (KP) postulates iff there exists a syncretic assignment which associates to every belief profile E a total pre-order \leq_E such that the result of the merging operation $\Delta_{IC}(E)$ as the set of minimal elements of $\text{Mod}(IC)$ according to the pre-order \leq_E .

Theorem

An operator Δ is a majority (resp. arbitration) merging operation iff there exists a majority (resp. fair) syncretic assignment which associates to every belief profile E a total pre-order \leq_E such that the result of the merging operation $\Delta_{IC}(E)$ as the set of minimal elements of $\text{Mod}(IC)$ according to the pre-order \leq_E .

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Distances

$d : \Omega \times \Omega \rightarrow \mathbb{N}$ is a distance between interpretations iff it respects

1 $\forall \omega_1, \omega_2 \in \Omega, d(\omega_1, \omega_2) = d(\omega_2, \omega_1)$

2 $d(\omega_1, \omega_2) = 0$ iff $\omega_1 = \omega_2$

It induces the distance between an interpretation and a formula:

$$d(\omega, \varphi) = \min_{\omega' \models \varphi} d(\omega, \omega')$$

Aggregation function

$f : \mathbb{N}^n \rightarrow \mathbb{N}$ is an aggregation function iff it respects

1 f is non-decreasing in each argument;

2 $\forall (x_1, \dots, x_n), f(x_1, \dots, x_n) = 0$ iff $x_1 = \dots = x_n = 0$;

3 $\forall x_1, f(x_1) = x_1$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Some distance functions:

drastic $d_D(\omega_1, \omega_2) = 0$ if $\omega_1 = \omega_2$, 1 otherwise

Hamming $d_H(\omega_1, \omega_2) = |\{x \in \mathcal{L} \mid \omega_1(x) \neq \omega_2(x)\}|$

Some aggregation functions: max, sum and lex.

Lexicographic aggregation

Given two vectors of numbers $\vec{a} = (a_1, \dots, a_n)$ and $\vec{b} = (b_1, \dots, b_n)$. Let σ and σ' be two permutations on $\{1, \dots, n\}$ s.t. $\forall i, a_{\sigma(i)} \geq a_{\sigma(i+1)}$ and $b_{\sigma'(i)} \geq b_{\sigma'(i+1)}$.

$\vec{a} \leq_{lex} \vec{b}$ iff $\forall i, a_{\sigma(i)} = b_{\sigma'(i)}$ or $\exists i \geq 1$ s.t. $a_{\sigma(i)} < b_{\sigma'(i)}$ and $a_{\sigma(j)} = b_{\sigma'(j)}$ for all $1 \leq j < i$.

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Distance-based merging operators

d is a distance, f and g are aggregation functions,
 $E = \{K_1, \dots, K_n\}$ is belief profile and C is a formula:

$$\text{Mod}(\Delta_{IC}^{d,f,g}(E)) = \{\omega \in \text{Mod}(IC) \mid d(\omega, E) \text{ is minimal} \}$$

where

$$d(\omega, E) = g(d(\omega, K_1), \dots, d(\omega, K_n))$$

and for every $K_i = \{\varphi_{i,1}, \dots, \varphi_{i,n_i}\}$

$$d(\omega, K_i) = f(d(\omega, \varphi_{i,1}), \dots, d(\omega, \varphi_{i,n_i}))$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Example

$E = \{K_1, K_2, K_3, K_4\}$ under the integrity constraint $IC = \top$ where

$$K_1 = \{a \wedge b \wedge c, a \rightarrow \neg b\}$$

$$K_2 = \{a \wedge b\}$$

$$K_3 = \{\neg a \wedge \neg b, \neg b\}$$

$$K_4 = \{a, a \rightarrow b\}$$

$\Delta^{d_H, \text{sum}, \text{lex}}$ Operator.

	$a \wedge b \wedge c$	$a \rightarrow \neg b$	$a \vee b$	$\neg a \wedge \neg b$	$\neg b$	a	$a \rightarrow b$	K_1, K_2, K_3, K_4	E
000	3	0	2	0	0	1	0	3, 2, 0, 1	3210
001	2	0	2	0	0	1	0	2, 2, 0, 1	2210
010	2	0	1	1	1	1	0	2, 1, 2, 1	2211
011	1	0	1	1	1	1	0	1, 1, 2, 1	2111
100	2	0	1	1	0	0	1	2, 1, 1, 1	2111
101	1	0	1	1	0	0	1	1, 1, 1, 1	1111
110	1	1	0	2	1	0	0	2, 0, 3, 0	3200
111	0	1	0	2	1	0	0	1, 0, 3, 0	3100

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Complexity for d_D

f/g	max	sum	lex
max	BH_2	Θ_2^p	Θ_2^p
sum	Θ_2^p	Θ_2^p	Δ_2^p

Complexity for d_H

f/g	max	sum	lex
max	Θ_2^p	Θ_2^p	Δ_2^p
sum	Θ_2^p	Θ_2^p	Δ_2^p

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

3 steps :

- subset of formulas which restore consistency: Potential Removed Sets
- minimal subset of formulas which restore consistency: Removed Sets
- profile without these formulas: Removed Sets Fusion operation

$E = \{K_1, \dots, K_n\}$: a belief profile IC : constraints
s.t. $K_1 \sqcup \dots \sqcup K_n \sqcup IC$ is inconsistent.
 X : a subset of formulas from $K_1 \sqcup \dots \sqcup K_n$.

Definition (Potential Removed Set)

X is a potential Removed Set of E constrained by IC iff
 $((K_1 \sqcup \dots \sqcup K_n) \setminus X) \sqcup IC$ is consistent.

$$K_1 = \{a \quad b\} \quad K_2 = \{\neg a \vee \neg b\}$$

Example

Potential Removed Sets

$$R_1 = \{a\}$$

$$R_2 = \{b\}$$

$$R_3 = \{\neg a \vee \neg b\}$$

$$R_4 = \{a \quad b\}$$

$$R_5 = \{b \quad \neg a \vee \neg b\}$$

$$R_6 = \{\neg a \vee \neg b \quad a\}$$

$$R_7 = \{\neg a \vee \neg b \quad a \quad b\}$$

Consistent subset

$$E \setminus R_1 = \{\neg a \vee \neg b \quad b\}$$

$$E \setminus R_2 = \{\neg a \vee \neg b \quad a\}$$

$$E \setminus R_3 = \{a \quad b\}$$

$$E \setminus R_4 = \{\neg a \vee \neg b\}$$

$$E \setminus R_5 = \{a\}$$

$$E \setminus R_6 = \{b\}$$

$$E \setminus R_7 = \emptyset$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

$E = \{K_1, \dots, K_n\}$: a belief profile IC : constraints
s.t. $K_1 \sqcup \dots \sqcup K_n \sqcup IC$ is inconsistent.
 P : a merging strategy.

Definition (Removed Set)

X is a Removed Set of E constrained by IC according to P iff :

- X is a potential Removed Set of E constrained by IC ;
- $\nexists X' \subseteq K_1 \sqcup \dots \sqcup K_n$ s.t. $X' \subset X$;
- $\nexists X' \subseteq K_1 \sqcup \dots \sqcup K_n$ s.t. $X' <_P X$.

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

$$K_1 = \{a \quad b\} \quad K_2 = \{\neg a \vee \neg b\}$$

Example

Removed Sets

$$R_1 = \{a\}$$

$$R_2 = \{b\}$$

$$R_3 = \{\neg a \vee \neg b\}$$

$$R_4 = \{a \quad b\}$$

$$R_5 = \{b \quad \neg a \vee \neg b\}$$

$$R_6 = \{\neg a \vee \neg b \quad a\}$$

$$R_7 = \{\neg a \vee \neg b \quad a \quad b\}$$

Consistent subset

$$E \setminus R_1 = \{\neg a \vee \neg b \quad b\}$$

$$E \setminus R_2 = \{\neg a \vee \neg b \quad a\}$$

$$E \setminus R_3 = \{a \quad b\}$$

$$E \setminus R_4 = \{\neg a \vee \neg b\}$$

$$E \setminus R_5 = \{a\}$$

$$E \setminus R_6 = \{b\}$$

$$E \setminus R_7 = \emptyset$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

Definition of the merging operator

$E = \{K_1, \dots, K_n\}$: a belief profile IC : constraints

P : a merging strategy.

$\mathcal{F}_{P,IC}\mathcal{R}(E)$: the set of Removed Sets of E constrained by IC according to P .

Definition ($\Delta_{P,IC}^{RSF}(E)$)

$$\Delta_{P,IC}^{RSF}(E) = \bigvee_{X \in \mathcal{F}_{P,IC}\mathcal{R}(E)} \{((K_1 \sqcup \dots \sqcup K_n) \setminus X) \sqcup IC\}$$

Example

$$K_1 = \{a \ b\} \quad K_2 = \{\neg a \vee \neg b\}$$

$$\Delta_{\Sigma,IC}^{RSF}(E) = \{\neg a \vee \neg b \ b\} \vee \{\neg a \vee \neg b \ a\} \vee \{a \ b\}$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

$E = \{K_1, \dots, K_n\}$: a belief profile.

X, X' : two potential Removed Sets of E .

Definition (\leq_Σ)

$$X \leq_\Sigma X' \text{ iff } \sum_{1 \leq i \leq n} |X \cap K_i| \leq \sum_{1 \leq i \leq n} |X' \cap K_i|$$

The *Sum* strategy



Profile $E = \{K_1, K_2, K_3\}$

$$K_1 = \{\neg d, \textcolor{green}{s} \vee o, \textcolor{green}{s}\} \quad K_2 = \{\neg s, d \vee o, \neg d \vee \neg o\}$$

$$K_3 = \{\textcolor{blue}{s}, \textcolor{blue}{d}, \textcolor{blue}{o}\}$$

6		$\textcolor{green}{s} \vee o$	$\textcolor{green}{s}$,	$d \vee o$,	$\textcolor{blue}{s}$,	$\textcolor{blue}{d}$,	$\textcolor{blue}{o}$
5		$\textcolor{green}{s} \vee o$,	$\neg \textcolor{green}{d}$,	$\textcolor{green}{s}$,	$\textcolor{blue}{s}$,	$\textcolor{blue}{o}$	
4		$\neg \textcolor{green}{d}$,	$\textcolor{green}{s}$,	$\neg d \vee \neg o$,	$\textcolor{blue}{s}$		
—		$d \vee o$,	$\neg s$,	$\textcolor{blue}{d}$,	$\textcolor{blue}{o}$		
3		$\textcolor{green}{s}$,	$\textcolor{green}{s}$,	$\textcolor{blue}{d}$			
—		$\neg \textcolor{green}{d}$,	$\neg s$,	$\textcolor{blue}{o}$			
—		$\neg \textcolor{green}{d}$,	$\neg s$,	$\neg d \vee \neg o$			
2		$\neg s$,	$\textcolor{blue}{d}$				

$$\Delta_{\Sigma, IC}^{RSF}(E) = \{\neg d \quad \textcolor{green}{s} \vee o \quad \textcolor{green}{s} \quad d \vee o \quad \neg d \vee \neg o \quad \textcolor{blue}{s} \quad \textcolor{blue}{o}\}$$

Introduction

Belief
revision

Several
sources -
belief
merging

Postulational
aspects

Distance-based
merging

Syntactic merging

Bibliography

4 Bibliography



**UNI
FREIBURG**

Introduction

Belief
revision

Several
sources -
belief
merging

Bibliography



Peter Gärdenfors and Hans Rott,
Belief revision,
Handbook of Logic in AI and LP, 1995.



Carlos E. Alchourrón, Peter Gärdenfors, David Makinson,
On the Logic of Theory Change: Partial Meet Contraction and Revision Functions,
Journal of Symbolic Logic, 1985.



Bernhard Nebel
Base Revision Operations and Schemes: Semantics, Representation and Complexity
ECAI, 1994.



Sébastien Konieczny, Jérôme Lang, Pierre Marquis:
DA2 merging operators
Artificial Intelligence, 2004.

Introduction

Belief
revision

Several
sources -
belief
merging

Bibliography



Julien Hué, Eric Würbel, Odile Papini:

Removed Sets Fusion: Performing Off The Shelf
ECAI, 2008.

Introduction

Belief
revision

Several
sources -
belief
merging

Bibliography