

Principles of Knowledge Representation and Reasoning

Qualitative Representation and Reasoning II: Allen's Interval Calculus

Bernhard Nebel, Malte Helmert and Stefan Wöfl

Albert-Ludwigs-Universität Freiburg

June 24, 2008

Principles of Knowledge Representation and Reasoning

June 24, 2008 — Qualitative Representation and Reasoning II: Allen's Interval Calculus
Allen's Interval Calculus

Motivation

Intervals and Relations Between Them

Processing an Example

Composition Table

Outlook

Reasoning in Allen's Interval Calculus

Constraint propagation algorithms (enforcing path consistency)

Example for Incompleteness

NP-Hardness Example

The Continuous Endpoint Class

Completeness for the CEP Class

A Maximal Tractable Sub-Algebra

The Endpoint Subclass

The ORD-Horn Subclass

Maximality

Solving Arbitrary Allen CSPs

Literature

Allen's Interval Calculus – Outline

Allen's Interval Calculus

Motivation

Intervals and Relations Between Them

Processing an Example

Composition Table

Outlook

Reasoning in Allen's Interval Calculus

A Maximal Tractable Sub-Algebra

Literature

Qualitative Temporal Representation and Reasoning

Often we do not want to talk about precise times:

- ▶ **NLP** – we do not have precise time points
- ▶ **Planning** – we do not want to commit to time points too early
- ▶ **Scenario descriptions** – we do not have the exact times or do not want to state them

What are the primitives in our representation system?

- ▶ **Time points**: actions and events are instantaneous, or we consider their beginning and ending
- ▶ **Time intervals**: actions and events have duration
- ▶ Reducibility? Expressiveness? Computational costs for reasoning?

Motivation: Example

Consider a planning scenario for multimedia generation:

- P1: *Display Picture1*
- P2: *Say "Put the plug in."*
- P3: *Say "The device should be shut off."*
- P4: *Point to Plug-in-Picture1.*

Temporal relations between events:

P2	should happen during	P1
P3	should happen during	P1
P2	should happen before or directly precede	P3
P4	should happen during or end together with	P2

- ↪ P4 happens before or directly precedes P3
- ↪ We could add the statement "P4 does not overlap with P3" without creating an inconsistency.

Allen's Interval Calculus

- ▶ Allen's interval calculus: **time intervals** and **binary relations** over them
- ▶ **Time intervals**: $X = (X^-, X^+)$, where X^- and X^+ are interpreted over the reals and $X^- < X^+$ (\rightsquigarrow naïve approach)
- ▶ **Relations** between concrete intervals, e. g.:
 - $(1.0, 2.0)$ strictly before $(3.0, 5.5)$
 - $(1.0, 3.0)$ meets $(3.0, 5.5)$
 - $(1.0, 4.0)$ overlaps $(3.0, 5.5)$
 - ...
- ↪ Which relations are conceivable?

The Base Relations


How many ways are there to order the four points of two intervals?

Relation	Symbol	Name
$\{(X, Y) : X^- < X^+ < Y^- < Y^+\}$	$<$	before
$\{(X, Y) : X^- < X^+ = Y^- < Y^+\}$	m	meets
$\{(X, Y) : X^- < Y^- < X^+ < Y^+\}$	o	overlaps
$\{(X, Y) : X^- = Y^- < X^+ < Y^+\}$	s	starts
$\{(X, Y) : Y^- < X^- < X^+ = Y^+\}$	f	finishes
$\{(X, Y) : Y^- < X^- < X^+ < Y^+\}$	d	during
$\{(X, Y) : Y^- = X^- < X^+ = Y^+\}$	\equiv	equal

and the **converse** relations (obtained by exchanging X and Y)

- ↪ These relations are JEPD.

The 13 Base Relations Graphically



Disjunctive Descriptions

- Assumption: We don't have precise information about the relation between X and Y , e. g.:

$$X \text{ o } Y \text{ or } X \text{ m } Y$$

- ... modelled by sets of base relations (meaning the union of the relations):

$$X \{o, m\} Y$$


$\rightsquigarrow 2^{13}$ imprecise relations (incl. \emptyset and \mathbf{B})

Example of an indefinite qualitative description:

$$\left\{ X \{o, m\} Y, Y \{m\} Z, X \{o, m\} Z \right\}$$

Our Example ... Formal

- P1: *Display* Picture1
- P2: *Say* "Put the plug in."
- P3: *Say* "The device should be shut off."
- P4: *Point* to Plug-in-Picture1.



Compose the constraints: $P4 \{d, f\} P2$ and $P2 \{d\} P1$: $P4 \{d\} P1$.

	λ	γ	δ	o	o ⁻¹	≡	≡ ⁻¹	s	s ⁻¹	f	f ⁻¹
λ	λ	λ	B	λ	λ	λ	λ	λ	λ	λ	λ
γ	B	γ	γ	γ	γ	γ	γ	γ	γ	γ	γ
δ	λ	γ	B	δ	δ	δ	δ	δ	δ	δ	δ
o	λ	γ	δ	B	o	o	o	o	o	o	o
o ⁻¹	λ	γ	δ	o	B	o ⁻¹	o ⁻¹	o ⁻¹	o ⁻¹	o ⁻¹	o ⁻¹
≡	λ	γ	δ	o	o ⁻¹	≡	≡	≡	≡	≡	≡
≡ ⁻¹	λ	γ	δ	o	o ⁻¹	≡	≡⁻¹	≡ ⁻¹	≡ ⁻¹	≡ ⁻¹	≡ ⁻¹
s	λ	γ	δ	o	o ⁻¹	≡	≡ ⁻¹	s	s	s	s
s ⁻¹	λ	γ	δ	o	o ⁻¹	≡	≡ ⁻¹	s	s⁻¹	s ⁻¹	s ⁻¹
f	λ	γ	δ	o	o ⁻¹	≡	≡ ⁻¹	s	s ⁻¹	f	f
f ⁻¹	λ	γ	δ	o	o ⁻¹	≡	≡ ⁻¹	s	s ⁻¹	f	f⁻¹

Outlook

- Using the **composition table** and the rules about operations on relations, we can **deduce** new relations between time intervals.
- What would be a **systematic** approach?
- How costly is that?
- Is that **complete**?
- If not, could it be complete on a subset of the relation system?

Reasoning in Allen's Interval Calculus

Allen's Interval Calculus

Reasoning in Allen's Interval Calculus

Constraint propagation algorithms (enforcing path consistency)

Example for Incompleteness

NP-Hardness Example

The Continuous Endpoint Class

Completeness for the CEP Class

A Maximal Tractable Sub-Algebra

Literature

Constraint Propagation – The Naive Algorithm


Enforcing path-consistency using the straight-forward method:

Let $Table[i, j]$ be an array of size $|n| \times |n|$ (n : number of intervals), in which we have recorded the constraints between the intervals.**EnforcePathConsistency1** (\mathcal{C}):*Input:* a (binary) CSP $\mathcal{C} = \langle V, D, C \rangle$ *Output:* an equivalent, but path consistent CSP \mathcal{C}' **repeat****for** each pair (i, j) , $1 \leq i, j \leq n$ **for** each k with $1 \leq k \leq n$ $Table[i, j] := Table[i, j] \cap (Table[i, k] \circ Table[k, j])$ **endfor****endfor****until** no entry in $Table$ is changed

↪ terminates;

↪ needs $O(n^5)$ intersections and compositions.An $O(n^3)$ Algorithm**EnforcePathConsistency2** (\mathcal{C}):*Input:* a (binary) CSP $\mathcal{C} = \langle V, D, C \rangle$ *Output:* an equivalent, but path consistent CSP \mathcal{C}' $Paths(i, j) = \{(i, j, k) : 1 \leq k \leq n\} \cup \{(k, i, j) : 1 \leq k \leq n\}$ $Queue := \bigcup_{i, j} Paths(i, j)$ **While** $Q \neq \emptyset$ select and delete (i, k, j) from Q $T := Table[i, j] \cap (Table[i, k] \circ Table[k, j])$ **if** $T \neq Table[i, j]$ $Table[i, j] := T$ $Table[j, i] := T^{-1}$ $Queue := Queue \cup Paths(i, j)$ **endif****endwhile**

Example for Incompleteness



NP-Hardness

Theorem (Kautz & Vilain)

CSAT is NP-hard for Allen's interval calculus.

Proof.

Reduction from 3-colorability (original proof using 3Sat).

Let $G = (V, E)$, $V = \{v_1, \dots, v_n\}$ be an instance of 3-colorability.

Then we use the intervals $\{v_1, \dots, v_n, 1, 2, 3\}$ with the following constraints:

$$\begin{array}{lll} 1 & \{m\} & 2 \\ 2 & \{m\} & 3 \\ v_i & \{m, \equiv, m^{-1}\} & 2 \quad \forall v_i \in V \\ v_i & \{m, m^{-1}, \prec, \succ\} & v_j \quad \forall (v_i, v_j) \in E \end{array}$$

This constraint system is satisfiable iff G can be colored with 3 colors. \square

Looking for Special Cases

- ▶ **Idea:** Let us look for polynomial special cases. In particular, let us look for sets of relations (subsets of the entire set of relations) that have an easy CSAT problem.
- ▶ **Note:** Interval formulae $X R Y$ can be expressed as clauses over atoms of the form $a \text{ op } b$, where:
 - ▶ a and b are endpoints X^-, X^+, Y^- and Y^+ and
 - ▶ $\text{op} \in \{<, >, =, \leq, \geq\}$.
- ▶ **Example:** All base relations can be expressed as unit clauses.

Lemma


Let $\pi(\Theta)$ be the translation of Θ to clause form. Θ is satisfiable over intervals iff $\pi(\Theta)$ is satisfiable over the rational numbers.

The Continuous Endpoint Class

Continuous Endpoint Class \mathcal{C} : This is a subset of \mathcal{A} such that there exists a clause form for each relation containing only unit clauses where $\neg(a = b)$ is forbidden.

Example: All basic relations and $\{d, o, s\}$, because

$$\pi(X \{d, o, s\} Y) = \begin{cases} X^- < X^+, Y^- < Y^+, \\ X^- < Y^+, X^+ > Y^-, \\ X^+ < Y^+ \end{cases}$$



Why Do We Have Completeness?

The set \mathcal{C} is closed under intersection, composition, and converse (it is a sub-algebra wrt. these three operations on relations). This can be shown by using a computer program.

Lemma

Each 3-consistent interval CSP over \mathcal{C} is globally consistent.

Theorem (van Beek)

Path consistency solves $\text{CMIN}(\mathcal{C})$ and decides $\text{CSAT}(\mathcal{C})$.

Proof.

Follows from the above lemma and the fact that a strongly n -consistent CSP is minimal. \square

Corollary

A path consistent interval CSP consisting of base relations only is satisfiable.

Helly's Theorem

Definition

A set $M \subseteq \mathbf{R}^n$ is **convex** iff for all pairs of points $a, b \in M$, all points on the line connecting a and b belong to M .

Theorem (Helly)

Let F be a family of at least $n + 1$ convex sets in \mathbf{R}^n . If all sub-families of F with $n + 1$ sets have a non-empty intersection, then $\bigcap F \neq \emptyset$.

Strong n -Consistency (1)

Proof.

We prove the claim by induction over k with $k \leq n$.

Base case: $k = 1, 2, 3$ \checkmark

Induction assumption: Assume strong $k - 1$ -consistency (and non-emptiness of all relations)

Induction step: From the assumption, it follows that there is an instantiation of $k - 1$ variables X_i to pairs (s_i, e_i) satisfying the constraints R_{ij} between the $k - 1$ variables.

We have to show that we can extend the instantiation to any k th variable.

Strong n -Consistency (2): Instantiating the k th Variable

Proof (Part 2).

The instantiation of the $k - 1$ variables X_i to (s_i, e_i) restricts the instantiation of X_k .

Note: Since $R_{ij} \in \mathcal{C}$ by assumption, these restrictions can be expressed by inequalities of the form:

$$s_i < X_k^+ \wedge e_j \geq X_k^- \wedge \dots$$

Such inequalities define convex subsets in \mathbf{R}^2 .

\rightsquigarrow Consider sets of 3 inequalities (= 3 convex sets).

Strong n -Consistency (3): Using Helly's Theorem

Proof (Part 3).

Case 1: All 3 inequalities mention only X_k^- (or mention only X_k^+). Then it suffices to consider only 2 of these inequalities (the strongest). Because of 3-consistency, there exists at least 1 common point satisfying these 3 inequalities.

Case 2: The inequalities mention X_k^- and X_k^+ , but it does not contain the inequality $X_k^- < X_k^+$. Then there are at most 2 inequalities with the same variable and we have the same situation as in Case 1.

Case 3: The set contains the inequality $X_k^- < X_k^+$. In this case, only three intervals (incl. X_k) can be involved and by the same argument as above there exists a common point.

\rightsquigarrow With Helly's Theorem, it follows that there exists a consistent instantiation for all subsets of variables.

\rightsquigarrow Strong k -consistency for all $k \leq n$.

Outlook

- ▶ $\text{CMIN}(\mathcal{C})$ can be computed in $O(n^3)$ time (for n being the number of intervals) using the path consistency algorithm.
- ▶ \mathcal{C} is a set of relations occurring “naturally” when observations are uncertain.
- ▶ \mathcal{C} contains 83 relations (incl. the impossible and the universal relations).
- ▶ Are there larger sets such that path consistency computes minimal CSPs? **Probably not**
- ▶ Are there larger sets of relations that permit polynomial satisfiability testing? **Yes**

A Maximal Tractable Sub-Algebra

Allen's Interval Calculus

Reasoning in Allen's Interval Calculus

A Maximal Tractable Sub-Algebra

The Endpoint Subclass

The ORD-Horn Subclass

Maximality

Solving Arbitrary Allen CSPs


Literature

The EP-Subclass

End-Point Subclass: $\mathcal{P} \subseteq \mathcal{A}$ is the subclass that permits a clause form containing only **unit** clauses ($a \neq b$ is allowed).

Example: all basic relations and $\{d, o\}$ since

$$\pi(X \{d, o\} Y) = \{X^- < X^+, Y^- < Y^+, \\ X^- < Y^+, X^+ > Y^-, X^- \neq Y^-, \\ X^+ < Y^+\}$$



Theorem (Vilain & Kautz 86, Ladkin & Maddux 88)

The path-consistency method decides $\text{CSAT}(\mathcal{P})$.

The ORD-Horn Subclass

ORD-Horn Subclass: $\mathcal{H} \subseteq \mathcal{A}$ is the subclass that permits a clause form containing only **Horn clauses**, where only the following **literals** are allowed:

$$a \leq b, a = b, a \neq b$$

$\neg a \leq b$ is not allowed!

Example: all $R \in \mathcal{P}$ and $\{o, s, f^{-1}\}$:

$$\pi(X \{o, s, f^{-1}\} Y) = \left\{ X^- \leq X^+, X^- \neq X^+, \\ Y^- \leq Y^+, Y^- \neq Y^+, \\ X^- \leq Y^-, \\ X^- \leq Y^+, X^- \neq Y^+, \\ Y^- \leq X^+, X^+ \neq Y^-, \\ X^+ \leq Y^+, \\ X^- \neq Y^- \vee X^+ \neq Y^+ \right\}.$$

Partial Orders: The ORD Theory

Let *ORD* be the following theory:

$$\begin{aligned} \forall x, y, z: \quad & x \leq y \wedge y \leq z \rightarrow x \leq z && \text{(transitivity)} \\ \forall x: \quad & x \leq x && \text{(reflexivity)} \\ \forall x, y: \quad & x \leq y \wedge y \leq x \rightarrow x = y && \text{(anti-symmetry)} \\ \forall x, y: \quad & x = y \rightarrow x \leq y && \text{(weakening of =)} \\ \forall x, y: \quad & x = y \rightarrow y \leq x && \text{(weakening of =)}. \end{aligned}$$

- ▶ *ORD* describes partially ordered sets, \leq being the ordering relation.
- ▶ *ORD* is a **Horn theory**
- ▶ What is missing wrt to *dense* and *linear* orders?

Satisfiability over Partial Orders

Proposition

Let Θ be a CSP over \mathcal{H} . Θ is satisfiable over interval interpretations iff $\pi(\Theta) \cup \text{ORD}$ is satisfiable over arbitrary interpretations.

Proof.

\Rightarrow : Since the reals form a partially ordered set (i. e., satisfy *ORD*), this direction is trivial.

\Leftarrow : Each extension of a partial order to a linear order satisfies all formulae of the form $a \leq b$, $a = b$, and $a \neq b$ which have been satisfied over the original partial order. \square

Complexity of CSAT(\mathcal{H})

Let $\text{ORD}_{\pi(\Theta)}$ be the propositional theory resulting from instantiating all axioms with the endpoints occurring in $\pi(\Theta)$.

Proposition

$\text{ORD} \cup \pi(\Theta)$ is satisfiable iff $\text{ORD}_{\pi(\Theta)} \cup \pi(\Theta)$ is so.

Proof idea: Herbrand expansion! \square

Theorem

$\text{CSAT}(\mathcal{H})$ can be decided in polynomial time.

Proof.

$\text{CSAT}(\mathcal{H})$ instances can be translated into a propositional Horn theory with blowup $O(n^3)$ according to the previous Prop., and such a theory is decidable in polynomial time. \square

$$\mathcal{C} \subset \mathcal{P} \subset \mathcal{H} \quad \text{with} \quad |\mathcal{C}| = 83, |\mathcal{P}| = 188, |\mathcal{H}| = 868$$

Path-Consistency and the OH-Class

Lemma

Let Θ be a path-consistent set over \mathcal{H} . Then

$$(X \{ \} Y) \notin \Theta \quad \text{iff} \quad \Theta \text{ is satisfiable}$$

Proof Idea.

One can show that $\text{ORD}_{\pi(\Theta)} \cup \pi(\Theta)$ is closed wrt **positive unit resolution**. Since this inference rule is refutation complete for Horn theories, the claim follows. \square

Lemma

\mathcal{H} is closed under intersection, composition, and conversion.

Theorem

The path-consistency method decides $\text{CSAT}(\mathcal{H})$.

\rightsquigarrow Maximality of \mathcal{H} ?

\rightsquigarrow Do we have to check all 8192 - 868 extensions?

Complexity of Sub-Algebras

Let \hat{S} be the closure of $S \subseteq \mathcal{A}$ under converse, intersection, and composition (i.e., the carrier of the least sub-algebra generated by S)

Theorem

$CSAT(\hat{S})$ can be polynomially transformed to $CSAT(S)$.

Proof Idea.

All relations in $\hat{S} - S$ can be modeled by a fixed number of compositions, intersections, and conversions of relations in S , introducing perhaps some fresh variables. \square

- \rightsquigarrow Polynomiality of S extends to \hat{S} .
- \rightsquigarrow NP-hardness of \hat{S} is inherited by all generating sets S .
- \rightsquigarrow Note: $\mathcal{H} = \hat{\mathcal{H}}$.

Minimal Extensions of the \mathcal{H} -Subclass

A computer-aided case analysis leads to the following result:

Lemma

There are only two minimal sub-algebras that strictly contain \mathcal{H} : $\mathcal{X}_1, \mathcal{X}_2$

$$N_1 = \{d, d^{-1}, o^{-1}, s^{-1}, f\} \in \mathcal{X}_1$$

$$N_2 = \{d^{-1}, o, o^{-1}, s^{-1}, f^{-1}\} \in \mathcal{X}_2$$

The clause form of these relations contain “proper” disjunctions!

Theorem

$CSAT(\mathcal{H} \cup \{N_i\})$ is NP-complete.

Question: Are there other maximal tractable subclasses?

“Interesting” Subclasses

Interesting subclasses of \mathcal{A} should contain all basic relations.

A computer-aided case analysis reveals: For $S \supseteq \{\{B\} : B \in \mathbf{B}\}$ it holds that

1. $\hat{S} \subseteq \mathcal{H}$, or
2. N_1 or N_2 is in \hat{S} .

In case 2, one can show: $CSAT(S)$ is NP-complete.

- \rightsquigarrow \mathcal{H} is the **only** maximal tractable subclass that is **interesting**.

Meanwhile, there is a **complete classification** of all sub-algebras containing at least one basic relation [IJCAI 2001] ... but the question for sub-algebras not containing a basic relation is open.

Relevance?

Theoretical:

We now know the boundary between polynomial and NP-hard reasoning problems along the dimension *expressiveness*. \oplus

Practical: All known applications either need only \mathcal{P} or they need more than \mathcal{H} ! \ominus

Backtracking methods might profit from the result because the branching factor is lower. $?$

- \rightsquigarrow How difficult is $CSAT(\mathcal{A})$ in practice?
- \rightsquigarrow What are the relevant branching factors?

Solving General Allen CSPs

- ▶ Backtracking algorithm using **path-consistency** as a **forward-checking method**
 - ▶ Relies on tractable fragments of Allen's calculus: split relations into relations of a tractable fragment, and backtrack over these.
 - ▶ Refinements and evaluation of different heuristics
- ↪ Which tractable fragment should one use?

Branching Factors

- ▶ If the labels are split into **base relations**, then on average a label is split into
6.5 relations
 - ▶ If the labels are split into **pointizable relations** (\mathcal{P}), then on average a label is split into
2.955 relations
 - ▶ If the labels are split into **ORD-Horn relations** (\mathcal{H}), then on average a label is split into
2.533 relations
- ↪ A difference of **0.422**
- ↪ This makes a difference for "hard" instances.

Summary

- ▶ Allen's interval calculus is often adequate for describing relative orders of events that have duration.
- ▶ The satisfiability problem for CSPs using the relations is NP-complete.
- ▶ For the **continuous endpoint class**, minimal CSPs can be computed using the path-consistency method.
- ▶ For the larger **ORD-Horn class**, CSAT is still decided by the path-consistency method.
- ▶ Can be used in practice for backtracking algorithms.

Literature I

- ▶ J.F. Allen.
Maintaining knowledge about temporal intervals.
Communications of the ACM, 26(11):832–843, November 1983.
Also in *Readings in Knowledge Representation*.
- ▶ P. van Beek and R. Cohen.
Exact and approximate reasoning about temporal relations.
Computational Intelligence, 6:132–144, 1990.
- ▶ B. Nebel and H.-J. Bürckert.
Reasoning about temporal relations: A maximal tractable subclass of Allen's interval algebra,
Journal of the ACM, 42(1): 43-66, 1995.
- ▶ B. Nebel.
Solving hard qualitative temporal reasoning problems: Evaluating the efficiency of using the ORD-horn class.
CONSTRAINTS, 1(3): 175-190, 1997.

Literature II

beamerr icon Article Krokhin, P. Jeavons and P. Jonsson.

A complete classification of complexity in Allen's algebra in the presence of a non-trivial basic relation.

Proc. 17th Int. Joint Conf. on AI (IJCAI-01), 83-88, Seattle, WA, 2001.