

Foundations of AI

13. Predicate Logic

Syntax and Semantics, Normal Forms,
Herbrand Expansion, Resolution

*Wolfram Burgard, Andreas Karwath,
Bernhard Nebel, and Martin Riedmiller*

13/1

Motivation

We can already do a lot with propositional logic. It is, however, annoying that there is no structure in the atomic propositions.

Example:

"All blocks are red"

"There is a block A"

It should follow that "A is red"

But propositional logic cannot handle this.

Idea: We introduce individual variables, predicates, functions, ...

→ First-Order Predicate Logic (PL1)

13/3

Contents

- Motivation
- Syntax and Semantics
- Normal Forms
- Reduction to Propositional Logic: Herbrand Expansion
- Resolution & Unification
- Closing Remarks

13/2

The Alphabet of First-Order Predicate Logic

Symbols:

- Operators: $\neg, \vee, \wedge, \forall, \exists, =$
- Variables: $x, x_1, x_2, \dots, x', x'', \dots, y, \dots, z, \dots$
- Brackets: $() , [] , \{ }$
- Function symbols (e.g., $weight(), color()$)
- Predicate symbols (e.g., $block(), red()$)
- Predicate and function symbols have an arity (number of arguments).
 - 0-ary predicate: propositional logic atoms
 - 0-ary function: constant
- We suppose a countable set of predicates and functions of any arity.
- "=" is usually not considered a predicate, but a logical symbol

13/4

The Grammar of First-Order Predicate Logic (1)

Terms (represent objects):

1. Every variable is a term.
2. If t_1, t_2, \dots, t_n are terms and f is an n-ary function, then

$$f(t_1, t_2, \dots, t_n)$$

is also a term.

Terms without variables: **ground terms**.

Atomic Formulae (represent statements about objects)

1. If t_1, t_2, \dots, t_n are terms and P is an n-ary predicate, then $P(t_1, t_2, \dots, t_n)$ is an atomic formula.
2. If t_1 and t_2 are terms, then $t_1 = t_2$ is an atomic formula.

Atomic formulae without variables: **ground atoms**. 13/5

Alternative Notation

Here	Elsewhere
$\neg\varphi$	$\sim\varphi \quad \bar{\varphi}$
$\varphi \wedge \psi$	$\varphi \& \psi \quad \varphi \bullet \psi \quad \varphi, \psi$
$\varphi \vee \psi$	$\varphi \mid \psi \quad \varphi; \psi \quad \varphi + \psi$
$\varphi \Rightarrow \psi$	$\varphi \rightarrow \psi \quad \varphi \supset \psi$
$\varphi \Leftrightarrow \psi$	$\varphi \leftrightarrow \psi \quad \varphi \equiv \psi$
$\forall x\varphi$	$(\forall x)\varphi \quad \wedge x\varphi$
$\exists x\varphi$	$(\exists x)\varphi \quad \vee x\varphi$

The Grammar of First-Order Predicate Logic (2)

Formulae:

1. Every atomic formula is a formula.
2. If φ and ψ are formulae and x is a variable, then
 - $\neg\varphi, \varphi \wedge \psi, \varphi \vee \psi, \varphi \Rightarrow \psi, \varphi \Leftrightarrow \psi, \exists x\varphi$ and $\forall x\varphi$ are also formulae.
 - \forall, \exists are as strongly binding as \neg .

Propositional logic is part of the PL1 language:

1. Atomic formulae: only 0-ary predicates
2. Neither variables nor quantifiers.

Meaning of PL1-Formulae

Our example: $\forall x[Block(x) \Rightarrow Red(x)], Block(a)$

For all objects x : If x is a block, then x is red and a is a block.

Generally:

- Terms are interpreted as objects.
- Universally-quantified variables denote all objects in the universe.
- Existentially-quantified variables represent one of the objects in the universe (made true by the quantified expression).
- Predicates represent subsets of the universe.

Similar to propositional logic, we define **interpretations, satisfiability, models, validity, ...**

Semantics of PL1-Logic

Interpretation: $I = \langle D, \bullet^I \rangle$ where D is an arbitrary, non-empty set and \bullet^I is a function that

- maps n -ary function symbols to functions over D :
 $f^I \in [D^n \rightarrow D]$
- maps individual constants to elements of D :
 $a^I \in D$
- maps n -ary predicate symbols to relations over D :
 $P^I \subseteq D^n$

Interpretation of ground terms:

$$(f(t_1, \dots, t_n))^I = f^I(t_1^I, \dots, t_n^I)$$

Satisfaction of ground atoms $P(t_1, \dots, t_n)$:

$$I \models P(t_1, \dots, t_n) \text{ iff } \langle t_1^I, \dots, t_n^I \rangle \in P^I$$

13/9

Example (1)

$$\begin{aligned} D &= \{d_1, \dots, d_n \mid n > 1\} \\ a^I &= d_1 \\ b^I &= d_2 \\ c^I &= \dots \\ \text{Block}^I &= \{d_1\} \\ \text{Red}^I &= D \\ I &\models \text{Red}(b) \\ I &\not\models \text{Block}(b) \end{aligned}$$

13/10

Example (2)

$$\begin{aligned} D &= \{1, 2, 3, \dots\} \\ 1^I &= 1 \\ 2^I &= 2 \\ &\dots \\ \text{Even}^I &= \{2, 4, 6, \dots\} \\ \text{succ}^I &= \{(1 \rightarrow 2), (2 \rightarrow 3), \dots\} \\ I &\models \text{Even}(2) \\ I &\not\models \text{Even}(\text{succ}(2)) \end{aligned}$$

13/11

Semantics of PL1: Variable Assignment

Set of all variables V . Function $\alpha: V \rightarrow D$

Notation: $\alpha[x/d]$ is the same as α apart from point x .

For x : $\alpha[x/d](x) = d$.

Interpretation of terms under I, α :

$$\begin{aligned} x^{I, \alpha} &= \alpha(x) \\ a^{I, \alpha} &= a^I \\ (f(t_1, \dots, t_n))^{I, \alpha} &= f^I(t_1^{I, \alpha}, \dots, t_n^{I, \alpha}) \end{aligned}$$

Satisfaction of atomic formulae:

$$I, \alpha \models P(t_1, \dots, t_n) \text{ iff } \langle t_1^{I, \alpha}, \dots, t_n^{I, \alpha} \rangle \in P^I$$

13/12

Example

$$\begin{aligned} \alpha &= \{(x \rightarrow d_1), (y \rightarrow d_2)\} \\ I, \alpha &\models Red(x) \\ I, \alpha[y/d_1] &\models Block(y) \end{aligned}$$

13/13

Example

$$\begin{aligned} T &= \{Block(a), Block(b), \forall x(Block(x) \Rightarrow Red(x))\} \\ D &= \{d_1, \dots, d_n \mid n > 1\} \\ a^I &= d_1 \\ b^I &= d_2 \\ Block^I &= \{d_1\} \\ Red^I &= D \\ \alpha &= \{(x \rightarrow d_1), (y \rightarrow d_2)\} \end{aligned}$$

Questions:

1. $I, \alpha \models Block(b) \vee \neg Block(b)$?
2. $I, \alpha \models Block(x) \Rightarrow (Block(x) \vee \neg Block(y))$?
3. $I, \alpha \models Block(a) \wedge Block(b)$?
4. $I, \alpha \models \forall x(Block(x) \Rightarrow Red(x))$?
5. $I, \alpha \models \top$?

13/15

Semantics of PL1: Satisfiability

A formula φ is satisfied by an interpretation I and a variable assignment α , i.e., $I, \alpha \models \varphi$:

$$\begin{aligned} I, \alpha &\models \top \\ I, \alpha &\not\models \perp \\ I, \alpha &\models \neg\varphi \text{ iff } I, \alpha \not\models \varphi \\ &\dots \end{aligned}$$

and all other propositional rules as well as

$$\begin{aligned} I, \alpha \models P(t_1, \dots, t_n) &\text{ iff } \langle t_1^{I, \alpha}, \dots, t_n^{I, \alpha} \rangle \in P^{I, \alpha} \\ I, \alpha \models \forall x\varphi &\text{ iff for all } d \in D, I, \alpha[x/d] \models \varphi \\ I, \alpha \models \exists x\varphi &\text{ iff there exists a } d \in D \text{ with } I, \alpha[x/d] \models \varphi \end{aligned}$$

13/14

Free and Bound Variables

$$\forall x[R(\boxed{y}, \boxed{z}) \wedge \exists y\{\neg P(y, x) \vee R(y, \boxed{z})\}]$$

The boxed appearances of y and z are **free**. All other appearances of x, y, z are **bound**.

Formulae with no free variables are called **closed** formulae or **sentences**. We form theories from closed formulae.

Note: With closed formulae, the concepts *logical equivalence, satisfiability, and implication, etc.* are not dependent on the variable assignment α (i.e., we can always ignore all variable assignments).

With closed formulae, α can be left out on the left side of the model relationship symbol:

$$I \models \varphi$$

13/16

Terminology

An interpretation I is called a **model** of φ under α if

$$I, \alpha \models \varphi$$

A PL1 formula φ can, as in propositional logic, be **satisfiable**, **unsatisfiable**, **falsifiable**, or **valid**.

Analogously, two formulae are **logically**

equivalent ($\varphi \equiv \psi$) if for all I, α :

$$I, \alpha \models \varphi \text{ iff } I, \alpha \models \psi$$

Note: $P(x) \not\equiv P(y)$!

Logical Implication is also analogous to propositional logic.

Question: How can we define **derivation**?

13/17

Equivalences for the Production of Prenex Normal Form

$$(\forall x\varphi) \wedge \psi \equiv \forall x(\varphi \wedge \psi) \text{ if } x \text{ not free in } \psi$$

$$(\forall x\varphi) \vee \psi \equiv \forall x(\varphi \vee \psi) \text{ if } x \text{ not free in } \psi$$

$$(\exists x\varphi) \wedge \psi \equiv \exists x(\varphi \wedge \psi) \text{ if } x \text{ not free in } \psi$$

$$(\exists x\varphi) \vee \psi \equiv \exists x(\varphi \vee \psi) \text{ if } x \text{ not free in } \psi$$

$$\forall x\varphi \wedge \forall x\psi \equiv \forall x(\varphi \wedge \psi)$$

$$\exists x\varphi \vee \exists x\psi \equiv \exists x(\varphi \vee \psi)$$

$$\neg \forall x\varphi \equiv \exists x\neg\varphi$$

$$\neg \exists x\varphi \equiv \forall x\neg\varphi$$

... and propositional logic equivalents

13/19

Prenex Normal Form

Because of the quantifiers, we cannot produce the CNF form of a formula directly.

First step: Produce the prenex normal form

quantifier prefix + (quantifier-free) matrix

$$Qx_1 Qx_2 Qx_3 \dots Qx_n \varphi$$

13/18

Production of Prenex Normal Form

1. Eliminate \Rightarrow and \Leftrightarrow
2. Move \neg inwards
3. Move quantifiers outwards

Example:

$$\begin{aligned} & \neg \forall x [(\forall x P(x)) \Rightarrow Q(x)] \\ \rightarrow & \neg \forall x [\neg(\forall x P(x)) \vee Q(x)] \\ \rightarrow & \exists x [(\forall x P(x)) \wedge \neg Q(x)] \end{aligned}$$

And now?

13/20

Renaming of Variables

$\varphi[\frac{x}{t}]$ is obtained from φ by replacing all free appearances of x in φ by t .

Lemma: Let y be a variable that does not appear in φ . Then it holds that

$$\forall x\varphi \equiv \forall y\varphi[\frac{x}{y}] \quad \text{and} \quad \exists x\varphi \equiv \exists y\varphi[\frac{x}{y}]$$

Theorem: There exists an algorithm that calculates the prenex normal form of any formula.

13/21

Skolemization

Idea: **Elimination of existential quantifiers** by applying a function that produces the "right" element.

Theorem (Skolem Normal Form): Let φ be a closed formula in prenex normal form such that all quantified variables are pair-wise distinct and the function symbols g_1, g_2, \dots do not appear in φ . Let

$$\varphi = \forall x_1 \dots \forall x_i \exists y \psi,$$

then φ is satisfiable iff

$$\varphi' = \forall x_1 \dots \forall x_i \psi[\frac{y}{g_i(x_1, \dots, x_i)}]$$

is satisfiable.

Example: $\forall x \exists y [P(x) \Rightarrow Q(y)] \rightarrow \forall x [P(x) \Rightarrow Q(g(x))]$

13/23

Derivation in PL1

Why is prenex normal form useful?

Unfortunately, there is no simple law as in propositional logic that allows us to determine satisfiability or general validity (by transformation into DNF or CNF).

But: we can **reduce the satisfiability problem in predicate logic to the satisfiability problem in propositional logic**. In general, however, this produces a very large number of propositional formulae (perhaps infinitely many)

Then: apply **resolution**.

13/22

Skolem Normal Form

Skolem Normal Form: Prenex normal form without existential quantifiers. Notation: φ^* is the SNF of φ .

Theorem: It is possible to calculate the Skolem normal form of every closed formula φ .

Example: $\exists x((\forall x P(x)) \wedge \neg Q(x))$ develops as follows:
 $\exists y((\forall x P(x)) \wedge \neg Q(y))$
 $\exists y(\forall x(P(x) \wedge \neg Q(y)))$
 $\forall x(P(x) \wedge \neg Q(g_0))$

Note: This transformation is **not an equivalence transformation**; it **only preserves satisfiability!**

Note: ... and is **not unique**.

13/24

Ground Terms & Herbrand Expansion

The set of **ground terms** (or **Herbrand Universe**) over a set of SNF formulae θ^* is the (infinite) set of all ground terms formed from the symbols of θ^* (in case there is no constant symbol, one is added). This set is denoted by $D(\theta^*)$.

The **Herbrand expansion** $E(\theta^*)$ is the instantiation of the Matrix ψ_i of all formulae in θ^* through all terms $t \in D(\theta^*)$:

$$E(\theta^*) = \{ \psi_i[\frac{x_1}{t_1}, \dots, \frac{x_n}{t_n}] \mid (\forall x_1, \dots, x_n \psi_i) \in \theta^*, t_j \in D(\theta^*) \}$$

Theorem (Herbrand): Let θ^* be a set of formulae in SNF. Then θ^* is satisfiable iff $E(\theta^*)$ is satisfiable.

Note: If $D(\theta^*)$ and θ^* are finite, then the Herbrand expansion is finite \rightarrow finite propositional logic theory.

Note: This is used heavily in AI and works well most of the time!

13/25

Recursive Enumeration and Decidability

We can construct a **semi-decision procedure** for **validity**, i.e., we can give a (rather inefficient) algorithm that *enumerates* all valid formulae step by step.

Theorem: The set of **valid** (and **unsatisfiable**) formulae in PL1 is **recursively enumerable**.

What about **satisfiable** formulae?

Theorem (undecidability of PL1): It is **undecidable**, whether a formula of PL1 is **valid**.

(Proof by reduction from PCP)

Corollary: The set of **satisfiable formulae** in PL1 is **not recursively enumerable**.

In other words: If a formula is valid, we can effectively confirm this fact. Otherwise, we can end up in an infinite loop.

13/27

Infinite Propositional Logic Theories

Can a **finite proof** exist when the set is infinite?

Theorem (compactness of propositional logic): A (countable) set of formulae of propositional logic is **satisfiable** if and only if **every finite subset is satisfiable**.

Corollary: A (countable) set of formulae in propositional logic is **unsatisfiable** if and only if a **finite subset is unsatisfiable**.

Corollary: (compactness of PL1): A (countable) set of formulae in predicate logic is **satisfiable** if and only if **every finite subset is satisfiable**.

13/26

Derivation in PL1

Clausal Form instead of Herbrand Expansion.

Clauses are **universally quantified disjunctions** of literals; all variables are universally quantified

$$\begin{array}{l} (\forall x_1, \dots, x_n)(l_1 \vee \dots \vee l_n) \text{ written as} \\ l_1 \vee \dots \vee l_n \quad \text{or} \\ \{l_1, \dots, l_n\} \end{array}$$

13/28

Production of Clausal Form from SNF

Skolem Normal Form

quantifier prefix + (quantifier-free) matrix

$$\forall x_1 \forall x_2 \forall x_3 \dots \forall x_n \varphi$$

1. Put Matrix into CNF using distribution rule
2. Eliminate universal quantifiers
3. Eliminate conjunction symbol
4. Rename variables so that no variable appears in more than one clause.

Theorem: It is possible to calculate the clausal form of every closed formula φ .

Note: Same remarks as for SNF

13/29

Conversion to CNF (1)

Everyone who loves all animals is loved by someone:

$$\forall x [\forall y \text{Animal}(y) \Rightarrow \text{Loves}(x, y)] \Rightarrow [\exists y \text{Loves}(y, x)]$$

1. Eliminate biconditionals and implications

$$\forall x \neg [\forall y \neg \text{Animal}(y) \vee \text{Loves}(x, y)] \vee [\exists y \text{Loves}(y, x)]$$

2. Move \neg inwards: $\neg \forall x p \equiv \exists x \neg p$, $\neg \exists x p \equiv \forall x \neg p$

$$\forall x [\exists y \neg (\neg \text{Animal}(y) \vee \text{Loves}(x, y))] \vee [\exists y \text{Loves}(y, x)]$$

$$\forall x [\exists y \neg \neg \text{Animal}(y) \wedge \neg \text{Loves}(x, y)] \vee [\exists y \text{Loves}(y, x)]$$

$$\forall x [\exists y \text{Animal}(y) \wedge \neg \text{Loves}(x, y)] \vee [\exists y \text{Loves}(y, x)]$$

13/30

Conversion to CNF (2)

3. Standardize variables: each quantifier should use a different one

$$\forall x [\exists y \text{Animal}(y) \wedge \neg \text{Loves}(x, y)] \vee [\exists z \text{Loves}(z, x)]$$

4. Skolemize: a more general form of existential instantiation. Each existential variable is replaced by a Skolem function of the enclosing universally quantified variables:

$$\forall x [\text{Animal}(F(x)) \wedge \neg \text{Loves}(x, F(x))] \vee [\text{Loves}(G(x), x)]$$

5. Drop universal quantifiers:

$$[\text{Animal}(F(x)) \wedge \neg \text{Loves}(x, F(x))] \vee [\text{Loves}(G(x), x)]$$

6. Distribute \wedge over \vee :

$$[\text{Animal}(F(x)) \vee \text{Loves}(G(x), x)] \wedge [\neg \text{Loves}(x, F(x)) \vee \text{Loves}(G(x), x)]$$

13/31

Clauses and Resolution

Assumption: All formulae in the KB are clauses.

Equivalently, we can assume that the KB is a *set of clauses*.

Due to commutativity, associativity, and idempotence of \vee , clauses can also be understood as *sets of literals*. The *empty set of literals* is denoted by \square .

Set of clauses: Δ

Set of literals: C, D

Literal: l

Negation of a literal: $\neg l$

13/32

Propositional Resolution

$$\frac{C_1 \cup \{l\}, C_2 \cup \{\bar{l}\}}{C_1 \cup C_2}$$

$C_1 \cup C_2$ are called **resolvents** of the **parent clauses** $C_1 \cup \{l\}$ and $C_1 \cup \{\bar{l}\}$. l and \bar{l} are the **resolution literals**.

Example: $\{a, b, \neg c\}$ resolves with $\{a, d, c\}$ to $\{a, b, d\}$.

Note: The resolvent is not equivalent to the parent clauses, but it follows from them!

Notation: $R(\Delta) = \Delta \cup \{C \mid C \text{ is a resolvent of two clauses from } \Delta\}$

13/33

What Changes?

Examples

$$\begin{aligned} & \{\{Nat(s(0)), \neg Nat(0)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\} \\ & \{\{Nat(s(0)), \neg Nat(x)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\} \\ & \{\{Nat(s(x)), \neg Nat(x)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\} \end{aligned}$$

We need **unification**, a way to make literals identical. Based on the notion of **substitution**, e.g., $\{\frac{x}{0}\}$.

13/34

Substitutions

A **substitution** $s = \{\frac{v_1}{t_1}, \dots, \frac{v_n}{t_n}\}$ **substitutes variables** v_i **for terms** t_i (t_i does NOT contain v_i).

Applying a substitution s to an expression φ yields the expression φs which is φ with all occurrences of v_i replaced by t_i for all i .

13/35

Substitution Examples

$$P(x, f(y), B)$$

$$P(z, f(w), B) \quad s = \{\frac{x}{z}, \frac{y}{w}\}$$

$$P(x, f(A), B) \quad s = \{\frac{y}{A}\}$$

$$P(g(z), f(A), B) \quad s = \{\frac{x}{g(z)}, \frac{y}{A}\}$$

$$P(C, f(A), A)$$

13/36

Composing Substitutions

Composing substitutions s_1 and s_2 gives s_1s_2 which is that substitution obtained by first applying to the terms in s_2 and adding s_1 remaining term/variable pairs (not occurring in s_2) to s_1 .

Example:
 $\{\frac{z}{g(x,y)}\} \{\frac{x}{A}, \frac{y}{B}, \frac{w}{C}, \frac{z}{D}\} = \{\frac{z}{g(A,B)}, \frac{x}{A}, \frac{y}{B}, \frac{w}{C}\}$

Application example:
 $P(x, y, z) \rightarrow P(A, B, g(A, B))$

13/37

Properties of substitutions

For a formula φ and substitutions s_1, s_2

$$(\varphi s_1) s_2 = \varphi (s_1 s_2)$$

$$(s_1 s_2) s_3 = s_1 (s_2 s_3) \quad \text{associativity}$$

$$s_1 s_2 \neq s_2 s_1 \quad \text{no commutativity!}$$

13/38

Unification

Unifying a set of expressions $\{w_i\}$

Find substitution s such that $w_i s = w_j s$ for all i, j

Example

$$\{P(x, f(y), B), P(x, f(B), B)\}$$

$$s = \{\frac{y}{B}, \frac{z}{A}\} \quad \text{not the simplest unifier}$$



$$s = \{\frac{y}{B}\} \quad \text{most general unifier (mgu)}$$

The **most general unifier**, the **mgu**, g of $\{w_i\}$ has the property that if s is any unifier of $\{w_i\}$ then there exists a substitution s' such that $\{w_i\} s = \{w_i\} g s'$

Property: The common instance produced is unique up to **alphabetic variants** (variable renaming)

13/39

Subsumption Lattice



13/40

Disagreement Set

The **disagreement set** of a set of expressions $\{w_i\}$ is the set of sub-terms $\{t_i\}$ of $\{w_i\}$ at the first position in $\{w_i\}$ for which the $\{w_i\}$ disagree

Examples

$\{P(x, A, f(y)), P(v, B, z)\}$ gives $\{x, v\}$
 $\{P(x, A, f(y)), P(x, B, z)\}$ gives $\{A, B\}$
 $\{P(x, y, f(y)), P(x, B, z)\}$ gives $\{y, B\}$

13/41

Example

$\{P(x, f(y), y), P(z, f(B), B)\}$

13/43

Unification Algorithm

Unify(Terms):

Initialize $k \leftarrow 0$;

Initialize $T_k = \text{Terms}$;

Initialize $s_k = \{\}$;

*If T_k is a singleton, then output s_k . Otherwise continue.

Let D_k be the disagreement set of T_k .

If there exists a var v_k and a term t_k in D_k such that v_k does not occur in t_k , continue. Otherwise, exit with failure.

$s_{k+1} \leftarrow s_k \{ \frac{v_k}{t_k} \}$;

$T_{k+1} \leftarrow T_k \{ \frac{v_k}{t_k} \}$;

$k \leftarrow k + 1$;

Goto *.

13/42

Binary Resolution

$$\frac{C_1 \cup \{l_1\}, C_2 \cup \{\bar{l}_2\}}{[C_1 \cup C_2]s}$$

where $s = \text{mgu}(l_1, \bar{l}_2)$, the most general unifier

$[C_1 \cup C_2]s$ is the **resolvent** of the **parent clauses** $C_1 \cup \{l_1\}$ and $C_2 \cup \{\bar{l}_2\}$.

$C_1 \cup \{l_1\}$ and $C_2 \cup \{\bar{l}_2\}$ do not share variables

l_1 and \bar{l}_2 are the **resolution literals**.

Examples: $\{\{Nat(s(0)), \neg Nat(0)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\}$
 $\{\{Nat(s(0)), \neg Nat(x)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\}$
 $\{\{Nat(s(x)), \neg Nat(x)\}, \{Nat(0)\}\} \vdash \{Nat(s(0))\}$

13/44

Some Further Examples

Resolve $P(x) \vee Q(f(x))$ and $R(g(x)) \vee \neg Q(f(A))$
 Standardizing the variables apart gives $P(x) \vee Q(f(x))$
 and $R(g(y)) \vee \neg Q(f(A))$
 Substitution $s = \{\frac{x}{A}\}$ Resolvent $P(A) \vee R(g(y))$

Resolve $P(x) \vee Q(x, y)$ and $\neg P(A) \vee \neg R(B, z)$
 Standardizing the variables apart
 Substitution $s = \{\frac{x}{A}\}$ and Resolvent $Q(A, y) \vee \neg R(B, z)$

13/45

Derivations

Notation: $R(\Delta) = \Delta \cup \{C \mid C \text{ is a resolvent or a factor of two clauses from } \Delta\}$

We say D can be **derived** from Δ , i.e.,

$$\Delta \vdash D,$$

If there exist $C_1, C_2, C_3, \dots, C_n = D$ such that

$C_i \in R(\Delta \cup \{C_1, \dots, C_{i-1}\})$ for $1 \leq i \leq n$.

13/47

Factoring

$$\frac{C_1 \cup \{l_1\} \cup \{l_2\}}{[CUl_1]s}$$

where $s = \text{mgu}(l_1, l_2)$ is the most general unifier.

Needed because:

$$\{\{P(u), P(v)\}, \{\neg P(x), \neg P(y)\}\} \models \square$$

but \square cannot be derived by binary resolution

Factoring yields:

$\{P(u)\}$ and $\{\neg P(x)\}$ whose resolvent is \square .

13/46

Example

From Russell and Norvig :

The law says it is a crime for an American to sell weapons to hostile nations. The country Nono, an enemy of America, has some missiles, and all of its missiles were sold to it by Colonel West, who is American.

Prove that Col. West is a criminal.

13/48

Example

... it is a crime for an American to sell weapons to hostile nations:

$American(x) \wedge weapon(y) \wedge Sells(x, y, z) \wedge Hostile(z) \Rightarrow Criminal(x)$

Nono ... has some missiles, i.e., $\exists x Ows(Nono, x) \wedge Missile(x)$:

$Ows(Nono, M_1)$ and $Missile(M_1)$

... all of its missiles were sold to it by Colonel West.

$\forall x Missiles(x) \wedge Ows(Nono, x) \Rightarrow Sells(West, x, Nono)$

Missiles are weapons:

$Missile(x) \Rightarrow Weapon(x)$

An enemy of America counts as "hostile":

$Enemy(x, America) \Rightarrow Hostile(x)$

West, who is American ...


$American(West)$

The country Nono, an enemy of America

$Enemy(Nono, America)$


13/49

An Example



13/50

Another Example



13/51

Properties of Resolution

Lemma: (soundness) If $\Delta \vdash D$, then $\Delta \models D$.

Lemma: resolution is refutation-complete:

Δ is unsatisfiable implies $\Delta \vdash \square$.

Theorem: Δ is unsatisfiable iff $\Delta \vdash \square$.

Technique:

to prove that $\Delta \models C$

negate C and prove that $\Delta \cup \{\neg C\} \vdash \square$.

13/52

The Lifting Lemma

Lemma:

Let C_1 and C_2 be two clauses with no shared variables, and let C_1' and C_2' be ground instances of C_1 and C_2 . If C' is a resolvent of C_1' and C_2' , then there exists a clause such that

- (1) C is a resolvent of C_1 and C_2
- (2) C' is a ground instance of C

Can be easily generalized to derivations

13/53

The general picture

Any set of sentences S is representable in clausal form

Assume S is unsatisfiable, and in clausal form

Some set S' of ground instances is unsatisfiable

Resolution can find a contradiction in S'

There is a resolution proof for the contradiction in S

← Herbrand's theorem

← Ground resolution theorem

← Lifting lemma

13/54

Closing Remarks: Processing

- **PL1-Resolution:** forms the basis of
 - most state of the art theorem provers for PL1
 - the programming language **Prolog**
 - only Horn clauses
 - considerably more efficient methods.
 - not dealt with : search/resolution strategies
- **Finite theories:** In applications, we often have to deal with a fixed set of objects. **Domain closure axiom:**
 - $\forall x[x = c_1 \vee x = c_2 \vee \dots \vee x = c_n]$
 - Translation into finite propositional theory is possible.

13/55

Closing Remarks: Possible Extensions

- PL1 is definitely very expressive, but in some circumstances we would like more ...
- **Second-Order Logic:** Also over predicate quantifiers
 - $\forall x, y[(x = y) \Leftrightarrow \{\forall p[p(x) \Leftrightarrow p(y)]\}]$
- Validity is no longer semi-decidable (we have lost compactness)
- **Lambda Calculus:** Definition of predicates, e.g., $\lambda x, y[\exists z P(x, z) \wedge Q(z, y)]$ defines a new predicate of arity 2
- Reducible to PL1 through Lambda-Reduction
- **Uniqueness quantifier:** $\exists! x \varphi(x)$ – there is exactly one x ...
- **Reduction to PL1:**
 - $\exists x[\varphi(x) \wedge \forall y\{\varphi(y) \Rightarrow x = y\}]$

13/56

Summary

- PL1 makes it possible to structure statements, thereby giving us considerably **more expressive power than propositional logic**.
- Formulae consist of **terms** and **atomic formulae**, which, together with **connectors** and **quantifiers**, can be put together to produce formulae.
- Interpretations in PL1 consist of a **universe** and an **interpretation function**.
- The **Herbrand Theory** shows that satisfiability in PL1 can be reduced to satisfiability in propositional logic (although infinite sets of formulae can arise under certain circumstances).
- **Resolution** is **refutation complete**
- **Validity** in PL1 is **not decidable** (it is only semi-decidable)