

Foundations of Artificial Intelligence

Prof. Dr. B. Nebel, Prof. Dr. M. Riedmiller
S. Lange, J. Witkowski, D. Zhang
Summer Term 2010

University of Freiburg
Department of Computer Science

Exercise Sheet 9

Due: Tuesday, July 6, 2010

Exercise 9.1 (Truth Tables, Models)

(a) Use truth tables to prove the validity of the following equivalences:

$$(i) (\alpha \rightarrow \beta) \equiv (\neg\alpha \vee \beta)$$

$$(ii) (\alpha \rightarrow \beta) \equiv (\neg\beta \rightarrow \neg\alpha)$$

$$(iii) \neg(\alpha \wedge \beta) \equiv (\neg\alpha \vee \neg\beta)$$

$$(iv) ((\alpha \wedge \beta) \vee \gamma) \equiv ((\alpha \vee \gamma) \wedge (\beta \vee \gamma))$$

(b) Consider a vocabulary with only four propositions, A , B , C , and D . How many models are there for the following formulae? Explain.

$$(i) (A \wedge B) \vee (B \wedge C)$$

$$(ii) A \vee B$$

$$(iii) (A \leftrightarrow B) \wedge (B \leftrightarrow C)$$

Exercise 9.2 (CNF Transformation, Resolution Method)

The following transformation rules hold, whereby propositional formulae can be transformed into equivalent formulae. Here, φ , ψ , and χ are arbitrary propositional formulae:

$$\neg\neg\varphi \equiv \varphi \tag{1}$$

$$\neg(\varphi \vee \psi) \equiv \neg\varphi \wedge \neg\psi \tag{2}$$

$$\varphi \vee (\psi \wedge \chi) \equiv (\varphi \vee \psi) \wedge (\varphi \vee \chi) \tag{3}$$

$$\neg(\varphi \wedge \psi) \equiv \neg\varphi \vee \neg\psi \tag{4}$$

$$\varphi \wedge (\psi \vee \chi) \equiv (\varphi \wedge \psi) \vee (\varphi \wedge \chi) \tag{5}$$

Additionally, the operators \vee and \wedge are associative and commutative.

Consider the formula $((C \wedge \neg B) \leftrightarrow A) \wedge (\neg C \rightarrow A)$.

(a) Transform the formula into a clause set K using the CNF transformation rules. Write down the steps.

(b) Afterwards, using the resolution method, show that $K \models (\neg B \rightarrow (A \wedge C))$ holds.

Exercise 9.3 (Functional Completeness, Negation Normal Form)

In this assignment, we only consider propositional logic *without* the implication and biimplication operators \rightarrow and \leftrightarrow . A propositional formula is in *Negation Normal Form* if negation symbols \neg only appear immediately in front of atoms.

- (a) Show: Each propositional formula can be transformed into an equivalent formula only containing the operators \neg and \vee .
- (b) Show: Each propositional formula can be transformed into an equivalent formula in Negation Normal Form.

The exercise sheets may and should be handed in and be worked on in groups of three (3) students. Please fill the cover sheet¹ and attach it to your solution.

¹<http://www.informatik.uni-freiburg.de/~ki/teaching/ss10/gki/coverSheet-english.pdf>